

## Optimierung

<http://www.math.uni-konstanz.de/numerik/personen/volkwein/teaching/>

### Sheet 1

**Deadline for hand-in: 2013/04/29 at lecture**

**Note:**

- Please write **each exercise** on a **separate sheet!**
- Remember to write **name, sheet number, exercise number** and your **group** on **each sheet!**

#### Exercise 1

Determine and identify the local critical point(s) of the Rosenbrock function

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x_1, x_2) = 100(x_2 - x_1^2)^2 + (1 - x_1)^2.$$

#### Exercise 2

Show that the function

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x_1, x_2) = 8x_1 + 12x_2 + x_1^2 - 2x_2^2$$

has only one stationary point, and that it is neither a maximum or minimum, but a saddle point. Sketch the contour lines of  $f$  (you can also use Matlab).

#### Exercise 3

(4 Points)

Consider the function

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x_1, x_2) = 3x_1^4 - 4x_1^2x_2 + x_2^2.$$

Prove that  $\tilde{x} = (0, 0)$  is a critical point of  $f$ . Show further, that a restriction of  $f$  on any line<sup>1</sup> through  $\tilde{x}$ , has a strict local minimum in  $\tilde{x}$ . Is  $\tilde{x}$  a local minimizer of  $f$ ?

---

<sup>1</sup>The restriction of  $f$  on the line  $\gamma$  is defined as  $g(t) = f(\gamma(t))$ ,  $t \in [0, 1]$ , with  $\gamma(t) := \tilde{x} + td$ , where  $d \in \mathbb{R}^2 \setminus \{0\}$  is an arbitrary but fixed direction.