## ÜBUNGEN ZU Theorie und Numerik partieller Differentialgleichungen

http://www.math.uni-konstanz.de/numerik/personen/volkwein/teaching/

## Sheet 1 Submission: 13.12.2010, 12:00 o'clock, Box 18

Exercise 1

Let  $\Omega \in \mathbb{R}^n$ ,  $n \in \mathbb{N}$ , be a connected and bounded domain with  $C^1$  boundary  $\partial\Omega$ . Let  $D_{\psi} = \{u \in C(\overline{\Omega}); u = \psi \text{ on } \partial\Omega\}$ . We consider the differential operator L:

$$L: C^2(\Omega) \cap D_0 \longrightarrow C(\Omega) , \quad Lu = -\Delta u.$$

Show:

i)  $\langle Lu, v \rangle = \langle u, Lv \rangle$  for  $u, v \in C^2(\Omega) \cap D_0$ ,

ii) L is positive definite.

The scalar product of two function u and v on  $\Omega$  is defined by

$$\langle u, v \rangle = \int_{\Omega} uv \, \mathrm{d}x \, .$$

## Exercise 2

Consider the linear problem

$$\begin{cases} -u''(x) + b(x)u'(x) + c(x)u(x) = f(x) \text{ for all } x \in (0,1), \\ u(0) = \alpha, u(1) = \beta, \end{cases}$$
(1)

where b, c, f are continuous functions on  $[0, 1] \subset \mathbb{R}$  with c > 0 and  $\alpha, \beta$  are real scalars. Discretize (3) using the step size  $h = \frac{1}{N}, N \in \mathbb{N}$ , and central/symmetric difference approximations for u''(x) und u'(x). Present the linear system. What are the conditions that the coefficient matrix of the obtained linear system is strictly diagonal dominant? What happens if u'(x) is approximated as follows:

$$u'(x) \approx \begin{cases} \frac{u(x+h) - u(x)}{h} & \text{if } b(x) < 0, \\ \frac{u(x) - u(x-h)}{h} & \text{if } b(x) \ge 0. \end{cases}$$
(2)

*Remark:* The approximation (2) is called *upwind scheme*. This kind of discretization is often used in the context of convection dominated equations.

(4 Points)

(4 Points)

*Hint:* Recall that a matrix  $A = ((a_{ij})) \in \mathbb{R}^{N \times N}$  is strictly row diagonal dominant if

$$\sum_{j=1, j \neq i}^{N} |a_{ij}| < |a_{ii}| \quad \text{for all } i \in \{1, \dots, N\}.$$

The matrix A is called row diagonal dominant if

$$\sum_{j=1, j \neq i}^{N} |a_{ij}| \le |a_{ii}| \quad \text{for all } i \in \{1, \dots, N\}$$

holds (see Lecture Notes Numerik I, page 16).

## Exercise 3

(4 Points)

We cosider the following elliptic problem in *divergence form*:

$$(a(x)v'(x))' = d(x) \quad \text{for } x \in \Omega = (0,1), \tag{3a}$$

where  $a \in C^1(\overline{\Omega})$  with  $a(x) \ge \underline{a} > 0$  for all  $x \in \Omega$  and  $d \in C(\Omega)$ . Together with (3a) we suppose Dirichlet boundary conditions

$$v(0) = \alpha, \quad v(1) = \beta \tag{3b}$$

with  $\alpha, \beta \in \mathbb{R}$ . The goal is to obtain a symmetric coefficient matrix when discritizing (3) by finite differences.

Discretize the interval  $\Omega$  using the equidistant grid  $x_i = ih$ ,  $0 \le i \le n+1$ , with the mesh size h = 1/(n+1),  $n \in \mathbb{N}$ . Approximate the outer derivative of (av')' at the grid point  $x_i$  by symmetric difference quotients using the intermediate grid points  $x_{i\pm 1/2} = x_i \pm h/2$ ,  $i = 1, \ldots, n$ . Write  $a_i = a(x_i)$ ,  $d_i = d(x_i)$  for  $i = 0, \ldots, n+1$  and  $a_{i\pm 1/2} = a(x_i \pm h/2)$  for  $i = 1, \ldots, n$ . Discretize the first derivative of v in the symmetric difference for (av')' by central differences. What are the equations that you derive for the approximation of (3)? Compose the coefficient matrix explicitly.