

c) Let  $\mathcal{C}(\mathbb{R})$  be the vector space of all real valued continuous functions on the real line. For any bounded interval  $[a, b]$  with  $a, b \in \mathbb{R}$  and  $a < b$ , we define for any  $f \in \mathcal{C}(\mathbb{R})$ :

$$p_{[a,b]}(f) := \sup_{a \leq t \leq b} |f(t)|.$$

$p_{[a,b]}$  is a seminorm but is never a norm because it might be that  $f(t) = 0$  for all  $t \in [a, b]$  (and so that  $p_{[a,b]}(f) = 0$ ) but  $f \neq 0$ . Other seminorms are the following ones:

$$q(f) := |f(0)| \quad \text{and} \quad q_p(f) := \left( \int_a^b |f(t)|^p dt \right)^{\frac{1}{p}} \quad \text{for } 1 \leq p < \infty.$$

Note that if  $0 < p < 1$  then  $q_p$  is not subadditive and so it is not a seminorm.

Seminorms on vector spaces are strongly related to a special kind of functionals, i.e. *Minkowski functionals*. Let us investigate more in details such a relation. Note that we are still in the realm of vector spaces with no topology!

**Definition 4.2.5.** Let  $X$  be a vector space and  $A$  a non-empty subset of  $X$ . We define the Minkowski functional (or gauge) of  $A$  to be the mapping:

$$\begin{aligned} p_A : X &\rightarrow \mathbb{R} \\ x &\mapsto p_A(x) := \inf\{\lambda > 0 : x \in \lambda A\} \end{aligned}$$

(where  $p_A(x) = \infty$  if the set  $\{\lambda > 0 : x \in \lambda A\}$  is empty).

It is then natural to ask whether there exists a class of subsets for which the associated Minkowski functionals are actually seminorms. The answer is positive for a class of subsets which we have already encountered in the previous section, namely for absorbing absolutely convex subsets. Actually we have even more as established in the following lemma.

**Notation 4.2.6.** Let  $X$  be a vector space and  $p$  a seminorm on  $X$ . The sets

$$\mathring{U}_p = \{x \in X : p(x) < 1\} \quad \text{and} \quad U_p = \{x \in X : p(x) \leq 1\}.$$

are said to be, respectively, the closed and the open unit semiball of  $p$ .

**Lemma 4.2.7.** Let  $X$  be a vector space. If  $A$  is a non-empty subset of  $X$  which is absorbing and absolutely convex, then the associated Minkowski functional  $p_A$  is a seminorm and  $\mathring{U}_{p_A} \subseteq A \subseteq U_{p_A}$ . Conversely, if  $q$  is a seminorm on  $X$  then  $\mathring{U}_q$  is an absorbing absolutely convex set and  $q = p_{\mathring{U}_q}$ .

*Proof.* Let  $A$  be a non-empty subset of  $X$  which is absorbing and absolutely convex and denote by  $p_A$  the associated Minkowski functional. We want to show that  $p_A$  is a seminorm.

- First of all, note that  $p_A(x) < \infty$  for all  $x \in X$  because  $A$  is absorbing. Indeed, by definition of absorbing set, for any  $x \in X$  there exists  $\rho_x > 0$  s.t. for all  $\lambda \in \mathbb{K}$  with  $|\lambda| \leq \rho_x$  we have  $\lambda x \in A$  and so the set  $\{\lambda > 0 : x \in \lambda A\}$  is never empty i.e.  $p_A$  has only finite nonnegative values. Moreover, since  $o \in A$ , we also have that  $o \in \lambda A$  for any  $\lambda \in \mathbb{K}$  and so  $p_A(o) = \inf\{\lambda > 0 : o \in \lambda A\} = 0$ .
- The balancedness of  $A$  implies that  $p_A$  is positively homogeneous. Since we have already showed that  $p_A(o) = 0$  it remains to prove the positive homogeneity of  $p_A$  for non-null scalars. Since  $A$  is balanced we have that for any  $x \in X$  and for any  $\xi, \lambda \in \mathbb{K}$  with  $\xi \neq 0$  the following holds:

$$\xi x \in \lambda A \text{ if and only if } x \in \frac{\lambda}{|\xi|} A. \quad (4.1)$$

Indeed,  $A$  balanced guarantees that  $\xi A = |\xi|A$  and so  $x \in \frac{\lambda}{|\xi|} A$  is equivalent to  $\xi x \in \lambda \frac{\xi}{|\xi|} A = \lambda A$ . Using (4.1), we get that for any  $x \in X$  and for any  $\xi \in \mathbb{K}$  with  $\xi \neq 0$ :

$$\begin{aligned} p_A(\xi x) &= \inf\{\lambda > 0 : \xi x \in \lambda A\} \\ &= \inf\left\{\lambda > 0 : x \in \frac{\lambda}{|\xi|} A\right\} \\ &= \inf\left\{|\xi| \frac{\lambda}{|\xi|} > 0 : x \in \frac{\lambda}{|\xi|} A\right\} \\ &= |\xi| \inf\{\mu > 0 : x \in \mu A\} = |\xi| p_A(x). \end{aligned}$$

- The convexity of  $A$  ensures the subadditivity of  $p_A$ . Take  $x, y \in X$ . By definition of Minkowski functional, for every  $\varepsilon > 0$  there exists  $\lambda, \mu > 0$  s.t.

$$\lambda \leq p_A(x) + \varepsilon \text{ and } x \in \lambda A$$

and

$$\mu \leq p_A(y) + \varepsilon \text{ and } y \in \mu A.$$

Then, by the convexity of  $A$ , we obtain that  $\frac{\lambda}{\lambda+\mu} A + \frac{\mu}{\lambda+\mu} A \subseteq A$ , i.e.  $\lambda A + \mu A \subseteq (\lambda + \mu)A$ , and therefore  $x + y \in (\lambda + \mu)A$ . Hence:

$$p_A(x + y) = \inf\{\delta > 0 : x + y \in \delta A\} \leq \lambda + \mu \leq p_A(x) + p_A(y) + 2\varepsilon$$

which proves the subadditivity of  $p_A$  since  $\varepsilon$  is arbitrary.

We can then conclude that  $p_A$  is a seminorm. Furthermore, we have the following inclusions:

$$\mathring{U}_{p_A} \subseteq A \subseteq U_{p_A}.$$

In fact, if  $x \in \mathring{U}_{p_A}$  then  $p_A(x) < 1$  and so there exists  $0 \leq \lambda < 1$  s.t.  $x \in \lambda A$ . Since  $A$  is balanced, for such  $\lambda$  we have  $\lambda A \subseteq A$  and therefore  $x \in A$ . On the other hand, if  $x \in A$  then clearly  $1 \in \{\lambda > 0 : x \in \lambda A\}$  which gives  $p_A(x) \leq 1$  and so  $x \in U_{p_A}$ .

Conversely, let us take any seminorm  $q$  on  $X$ . Let us first show that  $\mathring{U}_q$  is absorbing and absolutely convex and then that  $q$  coincides with the Minkowski functional associated to  $\mathring{U}_q$ .

- $\mathring{U}_q$  is absorbing.

Let  $x$  be any point in  $X$ . If  $q(x) = 0$  then clearly  $x \in \mathring{U}_q$ . If  $q(x) > 0$ , we can take  $0 < \rho < \frac{1}{q(x)}$  and then for any  $\lambda \in \mathbb{K}$  s.t.  $|\lambda| \leq \rho$  the positive homogeneity of  $q$  implies that  $q(\lambda x) = |\lambda|q(x) \leq \rho q(x) < 1$ , i.e.  $\lambda x \in \mathring{U}_q$ .

- $\mathring{U}_q$  is balanced.

For any  $x \in \mathring{U}_q$  and for any  $\lambda \in \mathbb{K}$  with  $|\lambda| \leq 1$ , again by the positive homogeneity of  $q$ , we get:  $q(\lambda x) = |\lambda|q(x) \leq q(x) < 1$  i.e.  $\lambda x \in \mathring{U}_q$ .

- $\mathring{U}_q$  is convex.

For any  $x, y \in \mathring{U}_q$  and any  $t \in [0, 1]$ , by both the properties of seminorm, we have that  $q(tx + (1-t)y) \leq tq(x) + (1-t)q(y) < t + 1 - t = 1$  i.e.  $tx + (1-t)y \in \mathring{U}_q$ .

Moreover, for any  $x \in X$  we easily see that

$$p_{\mathring{U}_q}(x) = \inf\{\lambda > 0 : x \in \lambda \mathring{U}_q\} = \inf\{\lambda > 0 : q(x) < \lambda\} = q(x). \quad \square$$

We are now ready to see the connection between seminorms and locally convex t.v.s..

**Definition 4.2.8.** *Let  $X$  be a vector space and  $\mathcal{P} := \{p_i\}_{i \in I}$  a family of seminorms on  $X$ . The coarsest topology  $\tau_{\mathcal{P}}$  on  $X$  s.t. each  $p_i$  is continuous is said to be the topology induced or generated by the family of seminorms  $\mathcal{P}$ .*

**Theorem 4.2.9.** *Let  $X$  be a vector space and  $\mathcal{P} := \{p_i\}_{i \in I}$  a family of seminorms. Then the topology induced by the family  $\mathcal{P}$  is the unique topology making  $X$  into a locally convex t.v.s. and having as a basis of neighbourhoods of the origin in  $X$  the following collection:*

$$\mathcal{B} := \left\{ \{x \in X : p_{i_1}(x) < \varepsilon, \dots, p_{i_n}(x) < \varepsilon\} : i_1, \dots, i_n \in I, n \in \mathbb{N}, \varepsilon > 0, \varepsilon \in \mathbb{R} \right\}.$$

*Viceversa, the topology of an arbitrary locally convex t.v.s. is always induced by a family of seminorms (often called generating).*

*Proof.* Let us first show that the collection  $\mathcal{B}$  is a basis of neighbourhoods of the origin for the unique topology  $\tau$  making  $X$  into a locally convex t.v.s. by using Theorem 4.1.14 and then let us prove that  $\tau$  actually coincides with the topology induced by the family  $\mathcal{P}$ .

For any  $i \in I$  and any  $\varepsilon > 0$ , consider the set  $\{x \in X : p_i(x) < \varepsilon\} = \varepsilon \mathring{U}_{p_i}$ . This is absorbing and absolutely convex, since we have already showed above that  $\mathring{U}_{p_i}$  fulfills such properties. Therefore, any element of  $\mathcal{B}$  is an absorbing absolutely convex subset of  $X$  as finite intersection of absorbing absolutely convex sets. Moreover, both properties a) and b) of Theorem 4.1.14 are clearly satisfied by  $\mathcal{B}$ . Hence, Theorem 4.1.14 guarantees that there exists a unique topology  $\tau$  on  $X$  s.t.  $(X, \tau)$  is a locally convex t.v.s. and  $\mathcal{B}$  is a basis of neighbourhoods of the origin for  $\tau$ .

Let us consider  $(X, \tau)$ . Then for any  $i \in I$ , the seminorm  $p_i$  is continuous, because for any  $\varepsilon > 0$  we have  $p_i^{-1}([0, \varepsilon]) = \{x \in X : p_i(x) < \varepsilon\} \in \mathcal{B}$  which means that  $p_i^{-1}([0, \varepsilon])$  is a neighbourhood of the origin in  $(X, \tau)$ . Therefore, the topology  $\tau_{\mathcal{P}}$  induced by the family  $\mathcal{P}$  is by definition coarser than  $\tau$ . On the other hand, each  $p_i$  is also continuous w.r.t.  $\tau_{\mathcal{P}}$  and so  $\mathcal{B} \subseteq \tau_{\mathcal{P}}$ . But  $\mathcal{B}$  is a basis for  $\tau$ , then necessarily  $\tau$  is coarser than  $\tau_{\mathcal{P}}$ . Hence,  $\tau \equiv \tau_{\mathcal{P}}$ .

Viceversa, let us assume that  $(X, \tau)$  is a locally convex t.v.s.. Then by Theorem 4.1.14 there exists a basis  $\mathcal{N}$  of neighbourhoods of the origin in  $X$  consisting of absorbing absolutely convex sets s.t. the properties a) and b) in Theorem 4.1.14 are fulfilled. W.l.o.g. we can assume that they are open. Consider now the family  $\mathcal{S} := \{p_N : N \in \mathcal{N}\}$ . By Lemma 4.2.7, we know that each  $p_N$  is a seminorm and that  $\mathring{U}_{p_N} \subseteq N$ . Let us show that for any  $N \in \mathcal{N}$  we have actually that  $N = \mathring{U}_{p_N}$ . Since any  $N \in \mathcal{N}$  is open and the scalar multiplication is continuous we have that for any  $x \in N$  there exists  $0 < t < 1$  s.t.  $x \in tN$  and so  $p_N(x) \leq t < 1$ , i.e.  $x \in \mathring{U}_{p_N}$ .

We want to show that the topology  $\tau_{\mathcal{S}}$  induced by the family  $\mathcal{S}$  coincides with original topology  $\tau$  on  $X$ . We know from the first part of the proof how to construct a basis for a topology induced by a family of seminorms. In fact, a basis of neighbourhoods of the origin for  $\tau_{\mathcal{S}}$  is given by

$$\mathcal{B} := \left\{ \bigcap_{i=1}^n \{x \in X : p_{N_i}(x) < \varepsilon\} : N_1, \dots, N_n \in \mathcal{N}, n \in \mathbb{N}, \varepsilon > 0, \varepsilon \in \mathbb{R} \right\}.$$

For any  $N \in \mathcal{N}$  we have showed that  $N = \mathring{U}_{p_N} \in \mathcal{B}$  so by Hausdorff criterion  $\tau \subseteq \tau_{\mathcal{S}}$ . Also for any  $B \in \mathcal{B}$  we have  $B = \bigcap_{i=1}^n \varepsilon \mathring{U}_{p_{N_i}} = \bigcap_{i=1}^n \varepsilon N_i$  for some  $n \in \mathbb{N}$ ,  $N_1, \dots, N_n \in \mathcal{N}$  and  $\varepsilon > 0$ . Then the property b) (of Theorem 4.1.14) for  $\mathcal{N}$  implies that for each  $i = 1, \dots, n$  there exists  $V_i \in \mathcal{N}$  s.t.  $V_i \subseteq \varepsilon N_i$  and so by the property a) of  $\mathcal{N}$  we have that there exists  $V \in \mathbb{N}$  s.t.  $V \subseteq \bigcap_{i=1}^n V_i \subseteq B$ . Hence, by Hausdorff criterion  $\tau_{\mathcal{S}} \subseteq \tau$ .  $\square$