

2.5 Completeness for t.v.s.

This section aims to treat completeness for most general types of topological vector spaces, beyond the traditional metric framework. As well as in the case of metric spaces, we need to introduce the definition of a Cauchy sequence in a t.v.s..

Definition 2.5.1. *A sequence $S := \{x_n\}_{n \in \mathbb{N}}$ of points in a t.v.s. X is said to be a Cauchy sequence if*

$$\forall U \in \mathcal{F}(o) \text{ in } X, \exists N \in \mathbb{N} : x_m - x_n \in U, \forall m, n \geq N. \quad (2.2)$$

This definition agrees with the usual one if the topology of X is defined by a translation-invariant metric d . Indeed, in this case, a basis of neighbourhoods of the origin is given by all the open balls centered at the origin. Therefore, $\{x_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence in such (X, d) iff $\forall \varepsilon > 0, \exists N \in \mathbb{N} : x_m - x_n \in B_\varepsilon(o), \forall m, n \geq N$, i.e. $d(x_m, x_n) = d(x_m - x_n, o) < \varepsilon$.

By using the subsequences $S_m := \{x_n \in S : n \geq m\}$ of S , we can easily rewrite (2.2) in the following way

$$\forall U \in \mathcal{F}(o) \text{ in } X, \exists N \in \mathbb{N} : S_N - S_N \subset U.$$

As we have already observed in Chapter 1, the collection $\mathcal{B} := \{S_m : m \in \mathbb{N}\}$ is a basis of the filter \mathcal{F}_S associated with the sequence S . This immediately suggests what the definition of a Cauchy filter should be:

Definition 2.5.2. *A filter \mathcal{F} on a subset A of a t.v.s. X is said to be a Cauchy filter if*

$$\forall U \in \mathcal{F}(o) \text{ in } X, \exists M \subset A : M \in \mathcal{F} \text{ and } M - M \subset U.$$

In order to better illustrate this definition, let us come back to our reference example of a t.v.s. X whose topology is defined by a translation-invariant metric d . For any subset M of (X, d) , recall that the diameter of M is defined as $\text{diam}(M) := \sup_{x, y \in M} d(x, y)$. Now if \mathcal{F} is a Cauchy filter on X then, by definition, for any $\varepsilon > 0$ there exists $M \in \mathcal{F}$ s.t. $M - M \subset B_\varepsilon(o)$ and this simply means that $\text{diam}(M) \leq \varepsilon$. Therefore, Definition 2.5.2 can be rephrased in this case as follows: *a filter \mathcal{F} on a subset A of such a metric t.v.s. X is a Cauchy filter if it contains subsets of A of arbitrarily small diameter.*

Going back to the general case, the following statement clearly holds.

Proposition 2.5.3. *The filter associated with a Cauchy sequence in a t.v.s. X is a Cauchy filter.*

Proposition 2.5.4.

Let X be a t.v.s.. Then the following properties hold:

- a) The filter of neighborhoods of a point $x \in X$ is a Cauchy filter on X .
- b) A filter finer than a Cauchy filter is a Cauchy filter.
- c) Every converging filter is a Cauchy filter.

Proof.

- a) Let $\mathcal{F}(x)$ be the filter of neighborhoods of a point $x \in X$ and let $U \in \mathcal{F}(o)$. By Theorem 2.1.10, there exists $V \in \mathcal{F}(o)$ such that $V - V \subset U$ and so such that $(V + x) - (V + x) \subset U$. Since X is a t.v.s., we know that $\mathcal{F}(x) = \mathcal{F}(o) + x$ and so $M := V + x \in \mathcal{F}(x)$. Hence, we have proved that for any $U \in \mathcal{F}(o)$ there exists $M \in \mathcal{F}(x)$ s.t. $M - M \subset U$, i.e. $\mathcal{F}(x)$ is a Cauchy filter.
- b) Let \mathcal{F} and \mathcal{F}' be two filters of subsets of X such that \mathcal{F} is a Cauchy filter and $\mathcal{F} \subseteq \mathcal{F}'$. Since \mathcal{F} is a Cauchy filter, by Definition 2.5.2, for any $U \in \mathcal{F}(o)$ there exists $M \in \mathcal{F}$ s.t. $M - M \subset U$. But \mathcal{F}' is finer than \mathcal{F} , so M belongs also to \mathcal{F}' . Hence, \mathcal{F}' is obviously a Cauchy filter.
- c) If a filter \mathcal{F} converges to a point $x \in X$ then $\mathcal{F}(x) \subseteq \mathcal{F}$ (see Definition 1.1.27). By a), $\mathcal{F}(x)$ is a Cauchy filter and so b) implies that \mathcal{F} itself is a Cauchy filter. □

The converse of c) is in general false, in other words not every Cauchy filter converges.

Definition 2.5.5. A subset A of a t.v.s. X is said to be complete if every Cauchy filter on A converges to a point x of A .

It is important to distinguish between completeness and sequentially completeness.

Definition 2.5.6. A subset A of a t.v.s. X is said to be sequentially complete if any Cauchy sequence in A converges to a point in A .

It is easy to see that complete always implies sequentially complete. The converse is in general false (see Example 2.5.9). We will encounter an important class of t.v.s., the so-called metrizable spaces, for which the two notions coincide.

Proposition 2.5.7. If a subset A of a t.v.s. X is complete then A is sequentially complete.

Proof.

Let $S := \{x_n\}_{n \in \mathbb{N}}$ a Cauchy sequence of points in A . Then Proposition 2.5.3 guarantees that the filter \mathcal{F}_S associated to S is a Cauchy filter in A . By the completeness of A we get that there exists $x \in A$ such that \mathcal{F}_S converges to x . This is equivalent to say that the sequence S is convergent to $x \in A$ (see Proposition 1.1.29). Hence, A is sequentially complete. \square

Before showing an example of a subset of a t.v.s. which is sequentially complete but not complete, let us introduce two useful properties about completeness in t.v.s..

Proposition 2.5.8.

- a) In a Hausdorff t.v.s. X , any complete subset is closed.
 b) In a complete t.v.s. X , any closed subset is complete.

Example 2.5.9.

Let $X := \mathbb{R}^d$ where $d > \aleph_0$ endowed with the product topology given by considering each copy of \mathbb{R} equipped with the usual topology given by the modulus. For convenience we write $X = \prod_{i \in J} \mathbb{R}$ with $|J| = d > \aleph_0$. Note that X is a Hausdorff t.v.s. as it is product of Hausdorff t.v.s.. Denote by H the subset of X consisting of all vectors $\underline{x} = (x_i)_{i \in J}$ in X with only countably many non-zero coordinates x_i . Claim: H is sequentially complete but not complete.

Proof. of Claim. Let us first make some observations on H .

- H is strictly contained in X .
 Indeed, any vector $\underline{y} \in X$ with all non-zero coordinates does not belong to H because $d > \aleph_0$.
- H is dense in X .
 In fact, let $\underline{x} = (x_i)_{i \in J} \in X$ and U a neighbourhood of \underline{x} in X . Then, by definition of product topology on X , there exist $\prod_{i \in J} U_i \subseteq U$ s.t. $U_i \in \mathbb{R}$ neighbourhood of x_i in \mathbb{R} for all $i \in J$ and $U_i \neq \mathbb{R}$ for all $i \in I$ where $I \subset J$ with $|I| < \infty$. Take $\underline{y} := (y_i)_{i \in J}$ s.t. $y_i \in U_i$ for all $i \in J$ with $y_i \neq 0$ for all $i \in I$ and $y_i = 0$ otherwise. Then clearly $\underline{y} \in U$ but also $\underline{y} \in H$ because it has only finitely many non-zero coordinates. Hence, $U \cap H \neq \emptyset$ and so $\overline{H} = X$.

Now suppose that H is complete, then by Proposition 2.5.8-a) we have that H is closed. Therefore, by the density of H in X , it follows that $H = \overline{H} = X$ which contradicts the first of the property above. Hence, H is not complete.

In the end, let us show that H is sequentially complete. Let $(\underline{x}_n)_{n \in \mathbb{N}}$ a Cauchy sequence of vectors $\underline{x}_n = (x_n^{(i)})_{i \in J}$ in H . Then for each $i \in J$ we have that the sequence of the i -th coordinates $(x_n^{(i)})_{n \in \mathbb{N}}$ is a Cauchy sequence

in \mathbb{R} . By the completeness (i.e. the sequentially completeness) of \mathbb{R} we have that for each $i \in J$, the sequence $(x_n^{(i)})_{n \in \mathbb{N}}$ converges to a point $x^{(i)} \in \mathbb{R}$. Set $\underline{x} := (x^{(i)})_{i \in J}$. Then:

- $\underline{x} \in H$, because for each $n \in \mathbb{N}$ only countably many $x_n^{(i)} \neq 0$ and so only countably many $x^{(i)} \neq 0$.
- the sequence $(\underline{x}_n)_{n \in \mathbb{N}}$ converges to \underline{x} in H . In fact, for any U neighbourhood of \underline{x} in X there exist $\prod_{i \in J} U_i \subseteq U$ s.t. $U_i \in \mathbb{R}$ neighbourhood of x_i in \mathbb{R} for all $i \in J$ and $U_i \neq \mathbb{R}$ for all $i \in I$ where $I \subset J$ with $|I| < \infty$. Since for each $i \in J$, the sequence $(x_n^{(i)})_{n \in \mathbb{N}}$ converges to $x^{(i)}$ in \mathbb{R} , we get that for each $i \in J$ there exists $N_i \in \mathbb{N}$ s.t. $x_n^{(i)} \in U_i$ for all $n \geq N_i$. Take $N := \max_{i \in I} N_i$ (the max exists because I is finite). Then for each $i \in J$ we get $x_n^{(i)} \in U_i$ for all $n \geq N$, i.e. $\underline{x}_n \in U$ for all $n \geq N$ which proves the convergence of $(\underline{x}_n)_{n \in \mathbb{N}}$ to \underline{x} .

Hence, we have showed that every Cauchy sequence in H is convergent. \square

In order to prove Proposition 2.5.8, we need two small lemmas regarding convergence of filters in a topological space.

Lemma 2.5.10. *Let \mathcal{F} be a filter of a topological Hausdorff space X . If \mathcal{F} converges to $x \in X$ and also to $y \in X$, then $x = y$.*

Proof.

Suppose that $x \neq y$. Then, since X is Hausdorff, there exists $V \in \mathcal{F}(x)$ and $W \in \mathcal{F}(y)$ such that $V \cap W = \emptyset$. On the other hand, we know by assumption that $\mathcal{F} \rightarrow x$ and $\mathcal{F} \rightarrow y$ that is $\mathcal{F}(x) \subseteq \mathcal{F}$ and $\mathcal{F}(y) \subseteq \mathcal{F}$ (see Definition 1.1.27). Hence, $V, W \in \mathcal{F}$. Since filters are closed under finite intersections, we get that $V \cap W \in \mathcal{F}$ and so $\emptyset \in \mathcal{F}$ which contradicts the fact that \mathcal{F} is a filter. \square

Lemma 2.5.11. *Let A be a subset of a topological space X . Then $x \in \overline{A}$ if and only if there exists a filter \mathcal{F} of subsets of X such that $A \in \mathcal{F}$ and \mathcal{F} converges to x .*

Proof.

Let $x \in \overline{A}$, i.e. for any $U \in \mathcal{F}(x)$ in X we have $U \cap A \neq \emptyset$. Set $\mathcal{F} := \{F \subseteq X \mid U \cap A \subseteq F \text{ for some } U \in \mathcal{F}(x)\}$. It is easy to see that \mathcal{F} is a filter of subsets of X . Therefore, for any $U \in \mathcal{F}(x)$, $U \cap A \in \mathcal{F}$ and $U \cap A \subseteq U$ imply that $U \in \mathcal{F}$, i.e. $\mathcal{F}(x) \subseteq \mathcal{F}$. Hence, $\mathcal{F} \rightarrow x$.

Viceversa, suppose that \mathcal{F} is a filter of X s.t. $A \in \mathcal{F}$ and \mathcal{F} converges to x . Let $U \in \mathcal{F}(x)$. Then $U \in \mathcal{F}$ since $\mathcal{F}(x) \subseteq \mathcal{F}$ by definition of convergence. Since also $A \in \mathcal{F}$ by assumption, we get $U \cap A \in \mathcal{F}$ and so $U \cap A \neq \emptyset$. \square