
Real Algebraic Geometry I – Exercise Sheet 9

Exercise 1 (4P). Give an algorithm deciding if a polynomial $f \in \mathbb{Q}[X, Y]$ has only finitely many roots in \mathbb{R}^2 .

Exercise 2 (4P). Let R be a real closed field and $n \in \mathbb{N}_0$. Show that a semialgebraic set $A \subseteq R^n$ has nonempty interior $\overset{\circ}{A}$ if and only if A is Zariski-dense in R^n (i.e., if no polynomial $f \in R[X_1, \dots, X_n] \setminus \{0\}$ vanishes on A).

Exercise 3 (8P). Show that the following sets are not K -semialgebraic:

- (a) the garden fence $\{(x, y) \in \mathbb{R}^2 \mid y \geq 0, y \leq \lfloor x \rfloor - x + \frac{1}{2}\} + 10\}$ where $K := \mathbb{R}$,
- (b) $\{(x, 2^x) \mid x \in \mathbb{R}\}$ where $K := \mathbb{Q}$,
- (c) the set of all *infinitesimal* elements in an arbitrary fixed non-archimedean real closed field R where $K := R$, and
- (d) the set $\{(x, y, z) \in R_{>0}^3 \mid \forall n \in \mathbb{N} : x \geq yn \geq zn^2\}$ for an arbitrary fixed non-archimedean real closed field R where $K := \mathbb{R}$.

Bonus exercise (4BP). Let R be a real closed field and A a finitely generated R -algebra. Suppose there exists an algebra homomorphism $\varphi: A \rightarrow S$ where S into a real closed extension field S of R .

- (a) Show that there exists also an algebra homomorphism $A \rightarrow R$.
- (b) Give a counterexample to (a) in the case where one drops the requirement that R is real closed.

Hint: For (a), find an ideal I with $\psi: A \xrightarrow{\cong} R[\underline{X}]/I$ and analyze the algebra homomorphism $\gamma: R[\underline{X}] \rightarrow R_1, f \mapsto \varphi(\psi^{-1}(\bar{f}^I))$ which is a point evaluation.

Please submit until Thursday, January 12, 2017, 11:44 in the box named RAG I, Number 10, near to the room F411.