

POSITIVE POLYNOMIALS LECTURE NOTES

(10: 18/05/10)

SALMA KUHLMANN

Contents

1. Ring of formal power series	1
2. Algebraic independence	4

1. RING OF FORMAL POWER SERIES

Definition 1.1. (Recall) Let $S = \{g_1, \dots, g_s\} \subseteq \mathbb{R}[x_1, \dots, x_n]$, then

$$\mathbf{K}_S := \{x \in \mathbb{R}^n \mid g_i(x) \geq 0 \forall i = 1, \dots, s\},$$

$\mathbf{T}_S := \left\{ \sum_{e_1, \dots, e_s \in \{0,1\}} \sigma_e g_1^{e_1} \dots g_s^{e_s} \mid \sigma_e \in \Sigma \mathbb{R}[\underline{X}]^2, e = (e_1, \dots, e_s) \right\}$ is the preordering generated by S .

Proposition 1.2. Let $n \geq 3$. Let S be a finite subset of $\mathbb{R}[\underline{x}]$ such that $K_S \subseteq \mathbb{R}^n$ has non empty interior. Then $\exists f \in \mathbb{R}[\underline{x}]$ such that $f \geq 0$ on \mathbb{R}^n and $f \notin T_S$.

To prove proposition 1.2 we need to learn a few facts about formal power series rings:

Definition 1.3. $\mathbb{R}[[\underline{x}]] := \mathbb{R}[[x_1, \dots, x_n]]$ **ring of formal power series** in $\underline{x} = (x_1, \dots, x_n)$ with coefficients in \mathbb{R} , i.e. , $f \in \mathbb{R}[[\underline{x}]]$ is expressible uniquely in the form

$$f = f_0 + f_1 + \dots,$$

where f_i is a homogenous polynomial of degree i in the variables x_1, \dots, x_n .

Here:

- Addition is defined point wise, and

- multiplication is defined using distributive law:

$$\left(\sum_{i=0}^{\infty} f_i\right)\left(\sum_{i=0}^{\infty} g_i\right) = (f_0g_0) + (f_0g_1 + f_1g_0) + (f_0g_2 + f_1g_1 + f_2g_0) + \dots = \sum_{k=0}^{\infty} \left(\sum_{i+j=k} f_i g_j\right)$$

So, both addition and multiplication are well defined and $\mathbb{R}[[x]]$ is an integral domain and $\mathbb{R}[x] \subseteq \mathbb{R}[[x]]$.

Notation 1.4. Fraction field of $\mathbb{R}[[x]]$ is denoted by

$$ff(\mathbb{R}[[x]]) := \mathbb{R}((x)).$$

The valuation $v : \mathbb{R}[[x]] \rightarrow \mathbb{Z} \cup \{\infty\}$ defined by:

$$v(f) = \begin{cases} \text{least } i \text{ s.t. } f_i \neq 0, & \text{if } f \neq 0 \\ \infty & \text{if } f = 0 \end{cases}$$

extends to $\mathbb{R}((x))$ via

$$v\left(\frac{f}{g}\right) := v(f) - v(g).$$

Lemma 1.5. Let $f \in \mathbb{R}[[x]]$; $f = f_k + f_{k+1} + \dots$, where f_i homogeneous of degree i , $f_k \neq 0$. Assume that f is a sos in $\mathbb{R}[[x]]$.

Then k is even and f_k is a sum of squares of forms of degree $\frac{k}{2}$.

Proof. $f = g_1^2 + \dots + g_l^2$, and

$$g_i = g_{ij} + g_{i,j+1} + \dots, \text{ with } j = \min\{v(g_i) ; i = 1, \dots, l\}$$

Then $f_0 = \dots = f_{2j-1} = 0$ and $f_{2j} = \sum_{i=1}^k g_{ij}^2 \neq 0$

So, $k = 2j$. □

1.6. Units in $\mathbb{R}[[x]]$: Let $f = f_0 + f_1 + \dots$, with $v(f) = 0$ i.e. $f_0 \neq 0$. Then f factors as

$$f = a(1 + t); \text{ where}$$

$$a \in R, a \neq 0, t \in \mathbb{R}[[x]] \text{ and } v(t) \geq 1 \text{ with } a := f_0 \in R \setminus \{0\}; t := \frac{1}{f_0}(f_1 + f_2 + \dots).$$

Lemma 1.7. $f \in \mathbb{R}[[x]]$ is a unit of $\mathbb{R}[[x]]$ if and only if $f_0 \neq 0$ (i.e. $v(f) = 0$).

Proof: $\frac{1}{1+t} = 1 - t + t^2 - \dots$, for $t \in \mathbb{R}[[x]] ; v(t) \geq 1$

is a well defined element of $\mathbb{R}[[\underline{x}]]$.

So, if $v(f) = 0$, then $f = a(1 + t)$ with $a \in \mathbb{R}, a \neq 0$ gives

$$f^{-1} = \frac{1}{a} \frac{1}{(1 + t)} \in \mathbb{R}[[\underline{x}]]. \quad \square$$

Corollary 1.8. It follows that $\mathbb{R}[[\underline{x}]]$ is a local ring because $I = \{f \mid v(f) \geq 1\}$ is a maximal ideal (quotient is a field \mathbb{R}).

Lemma 1.9. Let $f \in \mathbb{R}[[\underline{x}]]$ a positive unit, i.e. $f_0 > 0$. Then f is a square in $\mathbb{R}[[\underline{x}]]$.

Proof. $f = a(1 + t); a > 0, v(t) \geq 1$

$$\sqrt{f} = \sqrt{a} \sqrt{1 + t},$$

where $\sqrt{1 + t} := (1 + t)^{1/2} = 1 + \frac{1}{2}t - \frac{1}{8}t^2 + \dots$ is a well defined element of $\mathbb{R}[[\underline{x}]]$ □

Lemma 1.10. Suppose $n \geq 3$. Then $\exists f \in \mathbb{R}[\underline{x}]$ such that $f \geq 0$ on \mathbb{R}^n and f is not a sum of squares in $\mathbb{R}[\underline{x}]$.

Proof. Let $f \in \mathbb{R}[\underline{x}]$ be any homogeneous polynomial which is ≥ 0 on \mathbb{R}^n but is not a sum of squares in $\mathbb{R}[\underline{x}]$ (by Hilbert's Theorem such a polynomial exists).

Now by lemma 1.5 it follows that f is not sos in $\mathbb{R}[[\underline{x}]]$. □

Now we prove Proposition 1.2:

Proof of Proposition 1.2. Let $S = \{g_1, \dots, g_s\}$

• W.l.o.g. assume $g_i \neq 0$, for each $i = 1, \dots, s$. So $g := \prod_{i=1}^s g_i \neq 0$

$\text{int}(K_S) \neq \emptyset \Rightarrow \exists \underline{p} := (p_1, \dots, p_n) \in \text{int}(K_S)$ with $\prod_{i=1}^s g_i(\underline{p}) \neq 0$.

Thus $g_i(\underline{p}) > 0 \forall i = 1, \dots, s$.

• W.l.o.g. assume $\underline{p} = \underline{0}$ the origin

(by making a variable change $Y_i := X_i - p_i$, and noting that

$$\mathbb{R}[Y_1, \dots, Y_n] = \mathbb{R}[X_1, \dots, X_n])$$

So $g_i(0, \dots, 0) > 0$ for each $i = 1, \dots, s$ (i.e. has positive constant term),

that means $g_i \in \mathbb{R}[[\underline{X}]]$ is a positive unit in $\mathbb{R}[[\underline{X}]] \forall i = 1, \dots, s$.

By Lemma 1.9 (on positive units in power series): $g_i \in \mathbb{R}[[\underline{X}]]^2 \forall i = 1, \dots, s$.

So the preordering T_S^A generated by $S = \{g_1, \dots, g_s\}$ in the ring $A := \mathbb{R}[\underline{X}]$ is just $\Sigma\mathbb{R}[\underline{X}]^2$.

Now using Lemma 1.10 : $\exists f \in \mathbb{R}[\underline{X}]$, $f \geq 0$ on \mathbb{R}^n but f is not a sum of squares in $\mathbb{R}[\underline{X}]$ (i.e. $f \notin \Sigma\mathbb{R}[\underline{X}]^2 = T_S^{\mathbb{R}[\underline{X}]}$).

So clearly $f \notin T_S$. □(Proposition 1.2)

Proposition 1.2 that we just proved is just a special case of the following result due to Scheiderer:

Theorem 1.11. Let S be a finite subset of $\mathbb{R}[\underline{X}]$ such that K_S has dimension ≥ 3 . Then $\exists f \in \mathbb{R}[\underline{X}]$; $f \geq 0$ on \mathbb{R}^n and $f \notin T_S$.

To prove this result we need:

- (1) a reminder about dimension of semi algebraic sets, and
- (2) more facts about non singular zeros.

2. ALGEBRAIC INDEPENDENCE

Let E/F be a field extension:

Definition 2.1. (1) $a \in E$ is **algebraic** over F if it is a root of some non zero polynomial $f(x) \in F[x]$, otherwise a is a **transcendental** over F .

(2) $A = \{a_1, \dots, a_n\} \subseteq E$ is called **algebraically independent** over F if there is no nonzero polynomial $f(x_1, \dots, x_n) \in F[x_1, \dots, x_n]$ s.t. $f(a_1, \dots, a_n) = 0$.

In general $A \subseteq E$ is algebraically independent over F if every finite subset of A is algebraic independent over F .

(3) A **transcendence base** of E/F is a maximal subset (w.r.t. inclusion) of E which is algebraically independent over F .