

POSITIVE POLYNOMIALS LECTURE NOTES

(15: 08/06/10)

SALMA KUHLMANN

Contents

1. Schmüdgen's Nichtnegativstellensatz 1
2. Application of Schmüdgen's Positivstellensatz to the moment problem 2

1. SCHMÜDGEN'S NICHTNEGATIVSTELLENSATZ

1.1. Schmüdgen's Nichtnegativstellensatz (Recall 2.2.1 of lecture 14): Let K_S be a compact basic closed semi algebraic set and $f \in \mathbb{R}[X]$. Then

$$f \geq 0 \text{ on } K_S \Rightarrow \forall \epsilon \text{ real, } \epsilon > 0 : f + \epsilon \in T_S.$$

Corollary 1.2. Let $K = K_S$ be a compact basic closed semi algebraic set and $L \neq 0$ be a linear functional $L : \mathbb{R}[X] \rightarrow \mathbb{R}$ with $L(r) = r \forall r \in \mathbb{R}$. Then

$$\underbrace{L(T_S) \geq 0}_{\text{(i.e. } L(f) \geq 0 \forall f \in T_S)} \Rightarrow \underbrace{L(\text{Psd}(K_S)) \geq 0}_{\text{(i.e. } L(f) \geq 0 \forall f \geq 0 \text{ on } K_S)}.$$

Proof. W.l.o.g. $L(1) = 1, L \neq 0$. Let $f \in \text{Psd}(K_S)$ and assume $L(T_S) \geq 0$,

To show: $L(f) \geq 0$

By 1.1, $\forall \epsilon > 0 : f + \epsilon \in T_S$

So, $L(f + \epsilon) \geq 0$ i.e. $L(f) \geq -\epsilon \forall \epsilon > 0$ real

$\Rightarrow L(f) \geq 0$. □

We shall now relate this to the problem of representation of linear functionals via integration along measures (i.e. $\int d\mu$).

2. APPLICATION OF SPSS TO THE MOMENT PROBLEM

Let \mathcal{X} be a Hausdorff topological space.

Definition 2.1. \mathcal{X} is **locally compact** if $\forall x \in \mathcal{X} \exists$ open \mathcal{U} in \mathcal{X} s.t. $x \in \mathcal{U}$ and $\overline{\mathcal{U}}$ (closure) is compact.

Notation 2.2. $\mathcal{B}^\delta(\mathcal{X}) :=$ set of Borel measurable sets in \mathcal{X}
 = the smallest family of subsets of \mathcal{X} containing all compact subsets of \mathcal{X} , closed under finite \cup , set theoretic difference $A \setminus B$ and countable \cap .

Definition 2.3. A **Borel measure** μ on \mathcal{X} is a positive measure on \mathcal{X} s.t. every set in $\mathcal{B}^\delta(\mathcal{X})$ is measurable. We also require our measure to be **regular** i.e. $\forall B \in \mathcal{B}^\delta(\mathcal{X})$ and $\forall \epsilon > 0 \exists K, \mathcal{U} \in \mathcal{B}^\delta(\mathcal{X}), K$ compact, \mathcal{U} open s.t. $K \subseteq B \subseteq \mathcal{U}$ and $\mu(K) + \epsilon \geq \mu(B) \geq \mu(\mathcal{U}) - \epsilon$.

2.4. Moment problem is the following:

Given a closed set $K \subseteq \mathbb{R}^n$ and a linear functional $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$

Question:

$$\text{when does } \exists \text{ a Borel measure } \mu \text{ on } K \text{ s.t. } \forall f \in \mathbb{R}[\underline{X}] : L(f) = \int f d\mu ? \quad (1)$$

$$\text{Necessary condition for (1): } \forall f \in \mathbb{R}[\underline{X}], f \geq 0 \text{ on } K \Rightarrow L(f) \geq 0 \quad (2)$$

$$\text{in other words: } L(\text{Psd}(K)) \geq 0 \quad (3)$$

Is this necessary condition also sufficient?

The answer is YES.

Theorem 2.5. (Haviland) Given $K \subseteq \mathbb{R}^n$ closed and $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$ a linear functional with $L(1) > 0$:

$$\exists \mu \text{ as in (1) iff } \forall f \in \mathbb{R}[\underline{X}] : L(f) \geq 0 \text{ if } f \geq 0 \text{ on } K,$$

i.e. (1) \Leftrightarrow (2) \Leftrightarrow (3).

We shall prove Haviland's Theorem later. For now we shall deduce a corollary to SPSS.

Corollary 2.6. Let $K_S = \{x \mid g_i(x) \geq 0; i = 1, \dots, s\} \subseteq \mathbb{R}^n$ be a basic closed semi-algebraic set and compact, $L : \mathbb{R}[X] \rightarrow \mathbb{R}$ a linear functional with $L(1) > 0$. If $L(T_S) \geq 0$, then $\exists \mu$ positive Borel measure on K s.t. $L(f) = \int_{K_S} f d\mu \quad \forall f \in \mathbb{R}[X]$.

Remark 2.7. Let $S = \{g_1, \dots, g_s\}$.

1. $L(T_S) \geq 0$ can be written as

$$L(h^2 g_1^{e_1} \dots g_s^{e_s}) \geq 0 \quad \forall h \in \mathbb{R}[X], e_1, \dots, e_s \in \{0, 1\}.$$

2. Compare Haviland to Schmüdgen's moment problem, for compact K_S : we do not need to check $L(\text{Psd}(K_S)) \geq 0$ we only need to check $L(T_S) \geq 0$.

3. Reformulation of question (1) (in 2.4) in terms of moment sequences:

Let $L : \mathbb{R}[X] \rightarrow \mathbb{R}$, with $L(1) = 1$. Consider $\{X^\alpha = X_1^{\alpha_1} \dots X_n^{\alpha_n}; \alpha \in \mathbb{N}^n\}$ a monomial basis for $\mathbb{R}[X]$. So L is completely determined by the (multi)sequence of real numbers $\tau(\alpha) := L(X^\alpha)$; $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ (i.e. $\tau : \mathbb{N}^n \rightarrow \mathbb{R}$ is a function) and conversely, every such sequence determines a linear functional L :

$$L\left(\sum_{\alpha} a_{\alpha} X^{\alpha}\right) := \sum_{\alpha} a_{\alpha} L(X^{\alpha}).$$

So, (1) (in 2.4) can be reformulated as:

Given $K \subseteq \mathbb{R}^n$ closed, and a multisequence $\tau = \tau(\alpha)_{\alpha \in \mathbb{N}^n}$ of real numbers, $\exists \mu$ positive borel measure on K s.t $\int X^\alpha d\mu = \tau_\alpha$ for all $\alpha \in \mathbb{N}^n$?

Definition 2.8. A function $\tau : \mathbb{N}^n \rightarrow \mathbb{R}$ is a K -**moment sequence** if $\exists \mu$ positive borel measure on K s.t $\tau(\alpha) = \int_K X^\alpha d\mu$ for all $\alpha \in \mathbb{N}^n$

So (1) can be reformulated as: given K and a function $\tau : \mathbb{N}^n \rightarrow \mathbb{R}$, when is τ a K -moment sequence?

Definition 2.9. A function $\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$ is called **psd** if

$$\sum_{i,j=1}^m \tau(\underline{k}_i + \underline{k}_j) c_i c_j \geq 0,$$

for $m \geq 1$, arbitrary distinct $\underline{k}_1, \dots, \underline{k}_m \in (\mathbb{Z}_+)^n$; $c_1, \dots, c_m \in \mathbb{R}$.

Definition 2.10. Given $\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$ a function and a fixed polynomial

$g(\underline{X}) = \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} \underline{X}^{\underline{k}} \in \mathbb{R}[\underline{X}]$. Define a new function $g(E)_\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$ by

$g(E)_\tau(\underline{l}) := \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} \tau(\underline{k} + \underline{l})$; for any $\underline{l} \in (\mathbb{Z}_+)^n$.

Lemma 2.11. Let $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$ be a linear functional and denote by

$$\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$$

the corresponding multisequence (i.e. $\tau(\underline{k}) := L(\underline{X}^{\underline{k}}) \forall \underline{k} \in (\mathbb{Z}_+)^n$).

Fix $g \in \mathbb{R}[\underline{X}]$, $g(\underline{X}) = \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} \underline{X}^{\underline{k}} \in \mathbb{R}[\underline{X}]$. Then $L(h^2g) \geq 0$ for all $h \in \mathbb{R}[\underline{X}]$ if

and only if the multisequence $g(E)_\tau$ is psd.