

THE RING OF BOUNDED POLYNOMIALS ON A SEMI-ALGEBRAIC SET

DANIEL PLAUMANN AND CLAUD SCHEIDERER

ABSTRACT. Let V be a normal affine \mathbb{R} -variety, and let S be a semi-algebraic subset of $V(\mathbb{R})$ which is Zariski dense in V . We study the subring $B_V(S)$ of $\mathbb{R}[V]$ consisting of the polynomials that are bounded on S . We introduce the notion of S -compatible completions of V , and we prove the existence of such completions when $\dim(V) \leq 2$ or $S = V(\mathbb{R})$. An S -compatible completion X of V yields a ring isomorphism $\mathcal{O}_U(U) \xrightarrow{\sim} B_V(S)$ for some (concretely specified) open subvariety $U \supset V$ of X . We prove that $B_V(S)$ is a finitely generated \mathbb{R} -algebra if $\dim(V) \leq 2$ and S is open, and we show that this result becomes false in general when $\dim(V) \geq 3$.

INTRODUCTION

The general question studied in this paper can be stated as follows: Let S be a semi-algebraic subset of \mathbb{R}^n (i.e. a subset described by polynomial inequalities). How can we describe (conceptually or explicitly) the ring of polynomials that are bounded on S ?

To address this question we will work in the following setup. Let V be an affine variety defined over \mathbb{R} , and let S be a semi-algebraic subset of $V(\mathbb{R})$. We write $\mathbb{R}[V]$ for the ring of real polynomial functions on V (the coordinate ring of V) and consider its subring

$$B_V(S) = \{f \in \mathbb{R}[V] : f|_S \text{ is bounded}\}.$$

The first systematic study of $B_V(S)$, in the case $S = V(\mathbb{R})$, was undertaken in 1996 by Becker and Powers [2]. Their results have been generalized substantially by Monnier [12] and Schweighofer [20]. The emphasis there is on iterating the B_V construction (which requires a more abstract definition via the real spectrum), and on relations to sums of squares, sums of higher powers, and certificates for positivity; see also Marshall [11]. Motivation came in part from earlier work on the so-called holomorphy ring in the theory of real rings and fields (see [1] and [2] and references given there). In particular, rings A were studied in which all elements of the form $1 + \sum_i a_i^2$ with $a_i \in A$ are invertible. Of course, this condition is hardly ever satisfied for $A = \mathbb{R}[V]$.

A principal difficulty in studying $B_V(S)$ is that these rings need not be of finite type over \mathbb{R} . For example, this is so for elementary reasons when S is neither relatively compact nor Zariski dense in V (Cor. 5.8). We will show, however, that there exist other and more genuine examples as well (Cor. 5.14). In [2], [12] and [20],

2000 *Mathematics Subject Classification.* Primary 14P99; secondary 14C20, 14E15, 14P05.

Key words and phrases. Real algebraic varieties, bounded polynomials, normal varieties, divisors, finite generation of rings of global sections, algebraic surfaces.

this difficulty was overcome, to a certain extent, by working in the real spectrum and using sophisticated arguments from real algebra.

Our investigations go in a somewhat different direction. We seek to understand the structure of $B_V(S)$ in terms of the geometry of S , in particular by constructing algebraic compactifications of V that are suitably adapted to S . To describe the main results of this paper, let us make the simplifying assumption that the affine variety V is non-singular and connected. Using resolution of singularities, we construct a quasi-projective variety U containing V as an open dense subset such that $\mathcal{O}_U(U)$ restricts isomorphically onto $B_V(V(\mathbb{R}))$ (Thm. 3.8 and Thm. 4.1). We also generalize this construction from $S = V(\mathbb{R})$ to more general semi-algebraic sets $S \subset V(\mathbb{R})$. However we have been able to prove a satisfactory general result only for $\dim(V) \leq 2$ (Thm. 4.5). Combining this with a theorem of Zariski, we prove that the \mathbb{R} -algebra $B_V(S)$ is finitely generated when $\dim(V) \leq 2$ and the semi-algebraic set S satisfies a weak regularity condition (Thm. 5.12).

Here is a brief survey of the contents of this paper. After introducing the necessary terminology, we fix a connected normal affine \mathbb{R} -variety V and a closed semi-algebraic set $S \subset V(\mathbb{R})$, and we study morphisms $V \rightarrow W$ into affine \mathbb{R} -varieties W which are bounded on S . From this we obtain a characterization of the field of fractions of $B_V(S)$ in geometric terms (Prop. 2.2). In particular, we show that $B_V(S)$ has full transcendence degree $\dim(V)$ over \mathbb{R} if, and only if, there exists a non-constant $f \in B_V(S)$ such that the set $f^{-1}(c) \cap S$ is compact for almost all real numbers c (Thm. 2.5).

In Sect. 3 we introduce the notion of compatible completions. Given a normal affine variety V and a semi-algebraic set $S \subset V(\mathbb{R})$, a complete variety X containing V as an open dense set is said to be compatible with S if S touches every irreducible component of $X \setminus V$ in a Zariski dense subset, provided it touches that component at all. Removing those components from X that are not touched by S , we obtain an open subvariety $U \subset X$ containing V , and we show $\mathcal{O}_U(U) \xrightarrow{\sim} B_V(S)$ (Thm. 3.8). The existence of compatible completions is studied in Sect. 4. Relative to $S = V(\mathbb{R})$, every non-singular affine variety V has a compatible completion (Thm. 4.1). Relative to more general semi-algebraic subsets of $V(\mathbb{R})$, we can prove such a result for $\dim(V) \leq 2$ (Thm. 4.5). Turning things around, we prove in Thm. 4.11 that for every normal real quasi-projective variety U there is an open affine subset $V \subset U$ and a semi-algebraic set $S \subset V(\mathbb{R})$ with $\mathcal{O}_U(U) \xrightarrow{\sim} B_V(S)$. In Sect. 5 we finally study finite generation of the \mathbb{R} -algebra $B_V(S)$. After showing that $B_V(S)$ is not noetherian when S is not Zariski dense in V , we prove finite generation of $B_V(S)$ for $\dim(V) \leq 2$ (Thm. 5.12), using Zariski's theorem. We also show that this result becomes false when $\dim(V) \geq 3$.

Acknowledgements: Concerning the final section of this paper, we greatly profited from correspondence with Sebastian Krug, who provided us with examples and constructions in the context of Hilbert's fourteenth problem, going beyond what went into this paper. We are also grateful to Fabrizio Catanese for suggesting the connection to Zariski's work.

1. NOTATIONS AND CONVENTIONS

1.1. By an \mathbb{R} -variety we mean a separated and reduced \mathbb{R} -scheme V of finite type, not necessarily irreducible. The structural sheaf of V is written \mathcal{O}_V . If V is affine,

we write $\mathbb{R}[V] := \mathcal{O}_V(V)$ for the coordinate ring of V . If V is irreducible, then $\mathbb{R}(V)$ denotes the function field of V .

The set $V(\mathbb{R})$ of \mathbb{R} -rational points is endowed with the euclidean topology. The notion of semi-algebraic subsets of $V(\mathbb{R})$ is well-known when V is affine, and is easily transferred to the general case, a subset $S \subset V(\mathbb{R})$ being semi-algebraic if and only if $S \cap U(\mathbb{R})$ is semi-algebraic in $U(\mathbb{R})$ for every open affine subset U of V .

An irreducible \mathbb{R} -variety V is said to be *real* if it has a non-singular \mathbb{R} -point, or equivalently, if the function field $\mathbb{R}(V)$ is (formally) real. It is also equivalent that $V(\mathbb{R})$ is Zariski dense in V .

1.2. Let $f: V \rightarrow W$ be a morphism of \mathbb{R} -varieties, and let S be a subset of $V(\mathbb{R})$. We say that f is *bounded on S* if the closure of $f(S)$ in $W(\mathbb{R})$ is compact.

In particular, this applies in the case $W = \mathbb{A}^1$, i.e., when f is a regular function on V . The main object of this paper is to study the subring

$$B_V(S) := \{f \in \mathbb{R}[V]: f \text{ is bounded on } S\}$$

of $\mathbb{R}[V]$, for V an affine \mathbb{R} -variety. Here are some immediate observations:

Lemma 1.3. *Let V be an affine \mathbb{R} -variety, let S, S' be subsets of $V(\mathbb{R})$.*

- (a) $B_V(\overline{S}) = B_V(S)$;
- (b) $B_V(S) = \mathbb{R}[V]$ if and only if \overline{S} is compact;
- (c) $B_V(S \cup S') = B_V(S) \cap B_V(S')$;
- (d) if W is the (reduced) Zariski closure of S in V and I is the vanishing ideal of W in $\mathbb{R}[V]$, then $B_V(S)$ contains I , and $B_W(S) = B_V(S)/I$ as subrings of $\mathbb{R}[V]/I = \mathbb{R}[W]$;
- (e) the subring $B_V(S)$ is relatively integrally closed in $\mathbb{R}[V]$.

Proof. For (e), observe that a relation $f^n + a_1 f^{n-1} + \dots + a_n = 0$ with $f \in \mathbb{R}[V]$ and $a_1, \dots, a_n \in B_V(S)$ implies that $|f| \leq 1 + \max_i |a_i|$ holds (pointwise) on $V(\mathbb{R})$, hence f is bounded on S . \square

1.4. For the entire paper, our base field will be the field \mathbb{R} of real numbers. All results remain true when \mathbb{R} is replaced by an arbitrary real closed field and bounded is replaced by bounded over R , compact by semi-algebraically compact etc.

2. FIBRES OF BOUNDED MORPHISMS

Throughout this section we assume that V is an *irreducible* affine \mathbb{R} -variety and that S is a fixed semi-algebraic subset of $V(\mathbb{R})$.

2.1. Let $\varphi: V \rightarrow W$ be a morphism of \mathbb{R} -varieties. Given $y \in W(\mathbb{R})$, we write $S_y := \{x \in S: \varphi(x) = y\}$ for the fibre of y in S . Given $f \in \mathbb{R}[V]$, we define

$$\Omega_\varphi(f) := \{y \in W(\mathbb{R}): f \text{ is unbounded on } S_y\}.$$

This is a semi-algebraic subset of $W(\mathbb{R})$ which is contained in $\varphi(S)$.

Proposition 2.2. *Let $B = B_V(S)$. For $f \in \mathbb{R}[V]$, the following are equivalent:*

- (i) $f \in \text{Quot}(B)$;
- (ii) there exists a dominant morphism $\varphi: V \rightarrow W$ of affine \mathbb{R} -varieties which is bounded on S such that $\Omega_\varphi(f)$ is not Zariski dense in W .

Proof. When V is an affine \mathbb{R} -variety and $f \in \mathbb{R}[V]$, we write $Z(f) := \{x \in V(\mathbb{R}) : f(x) = 0\}$ for the zero set of f in $V(\mathbb{R})$. More generally, if $M \subset V(\mathbb{R})$ is a semi-algebraic subset and $f: M \rightarrow \mathbb{R}$ is a (semi-algebraic) map, we denote the zero set of f in M by $Z(f)$.

Assume $f \in \text{Quot}(B)$, so there is $0 \neq h \in B$ with $fh \in B$. Let C be any finitely generated subalgebra of B containing h and fh , write $W = \text{Spec}(C)$, and let $\varphi: V \rightarrow W$ be the morphism induced by the inclusion $C \subset \mathbb{R}[V]$. Then φ is bounded on S . Let $y \in \Omega_\varphi(f)$, so f is unbounded on $S_y = S \cap \varphi^{-1}(y)$. On the other hand, h and fh lie in $C = \mathbb{R}[W]$, so they are constant on S_y . Together, this implies $h(y) = 0$. So $\Omega_\varphi(f)$ is contained in $Z(h)$, and is therefore not Zariski dense in W .

Conversely assume that there is a dominant morphism $\varphi: V \rightarrow W$ as in (ii). Via φ , we consider $\mathbb{R}[W]$ as a subring of $B_V(S) \subset \mathbb{R}[V]$. Write $\Omega := \Omega_\varphi(f)$. We first define a map $\tilde{f}: \overline{\varphi(S)} \rightarrow \mathbb{R}$ by

$$\tilde{f}(y) := \begin{cases} \left(1 + \sup\{|f(x)| : x \in S_y\}\right)^{-1} & \text{if } S_y \neq \emptyset, \\ 1 & \text{if } S_y = \emptyset, \end{cases}$$

($y \in \overline{\varphi(S)}$), where we put $\frac{1}{\infty} := 0$. The map \tilde{f} has semi-algebraic graph, and $\tilde{f}^{-1}(0) = \Omega$. Let D be the set of points in $\overline{\varphi(S)}$ where \tilde{f} fails to be continuous. Then D is not Zariski dense in W . This can, for instance, easily be deduced from a cylindrical algebraic decomposition of the graph of \tilde{f} (see [3] Thm. 2.3.1). Hence $D \cup \Omega$ is not Zariski dense in W either.

By Lemma 2.3 below, there exists a continuous map $\tilde{h}: \overline{\varphi(S)} \rightarrow \mathbb{R}$ with semi-algebraic graph such that $|\tilde{h}| \leq \tilde{f}$ on $\overline{\varphi(S)}$, and such that $Z(\tilde{h}) \subset \overline{D} \cup \Omega$. From the definition of \tilde{f} it is clear that $|f(x)\tilde{h}(\varphi(x))| \leq |f(x)\tilde{f}(\varphi(x))| < 1$ for every $x \in S$.

Since $Z(\tilde{h})$ is not Zariski dense in W , there exists $h \neq 0$ in $\mathbb{R}[W]$ with $Z(\tilde{h}) \subset Z(h)$. Since $\overline{\varphi(S)}$ is compact, the Łojasiewicz inequality (see [3], Cor. 2.6.7) implies that there are $0 < c \in \mathbb{R}$ and a positive integer N such that $|h^N| \leq c \cdot |\tilde{h}|$ on $\overline{\varphi(S)}$. We conclude

$$|f(x)h(\varphi(x))^N| \leq c \cdot |f(x)\tilde{h}(\varphi(x))| < c$$

for all $x \in S$. This shows that fh^N is bounded on S , and so $f = (fh^N)/h^N$ lies in $\text{Quot}(B)$. \square

The following easy fact was used in the last proof:

Lemma 2.3. *Let $M \subset \mathbb{R}^n$ be a semi-algebraic subset, and let $f: M \rightarrow \mathbb{R}$ be a map with semi-algebraic graph and with $f \geq 0$ on M . Let D be the set of points in M where f fails to be continuous. Then there exists a continuous semi-algebraic function $g: M \rightarrow \mathbb{R}$ such that $|g| \leq f$ on M and such that $Z(g) \subset Z(f) \cup \overline{D}$.*

Proof. Upon replacing f by $\min\{f, 1\}$ we may assume $f \leq 1$ on M . The distance function $d_D: \mathbb{R}^n \rightarrow \mathbb{R}$, $d_D(x) = \inf\{|x - y| : y \in D\}$ is continuous with semi-algebraic graph, and it vanishes precisely on \overline{D} . The function

$$g(x) := f(x) \cdot \frac{d_D(x)}{1 + d_D(x)} \quad (x \in M)$$

has the desired properties. Indeed, f is continuous outside of \overline{D} , and hence so is g . Fix $x \in M \cap \overline{D}$. Then $g(x) = 0$, and for all $y \in M$ we have $d_D(y) \leq |y - x|$. This

implies

$$0 \leq g(y) \leq \frac{|y-x|}{1+|y-x|}$$

for all $y \in M$. In particular, g is continuous in x . \square

2.4. We fix an affine irreducible \mathbb{R} -variety V and a semi-algebraic set $S \subset V(\mathbb{R})$. Given a morphism $\varphi: V \rightarrow W$ of \mathbb{R} -varieties, we put

$$\Omega_\varphi := \bigcup_{f \in \mathbb{R}[V]} \Omega_\varphi(f).$$

If $\mathbb{R}[V]$ is generated by x_1, \dots, x_n as an \mathbb{R} -algebra then $\Omega_\varphi = \bigcup_{i=1}^n \Omega_\varphi(x_i)$. This shows that Ω_φ is a semi-algebraic subset of $W(\mathbb{R})$. Clearly,

$$\Omega_\varphi = \{y \in W(\mathbb{R}) : \overline{S_y} \text{ is not compact}\}.$$

Given an \mathbb{R} -algebra B , we define the transcendence degree $\text{trdeg}_{\mathbb{R}}(B)$ of B as the maximum number of elements of B which are algebraically independent over \mathbb{R} .

Theorem 2.5. *Let $S \subset V(\mathbb{R})$ be a semi-algebraic set, let $B = B_V(S)$. The following conditions are equivalent:*

- (i) $\text{Quot}(B) = \mathbb{R}(V)$;
- (ii) $\text{trdeg}_{\mathbb{R}}(B) = \dim(V)$;
- (iii) *there is a birational morphism $V \rightarrow W$ of affine varieties which is bounded on S ;*
- (iv) *there is a dominant morphism $\varphi: V \rightarrow W$ of affine varieties which is bounded on S such that Ω_φ is not Zariski dense in W ;*
- (v) *there is a non-constant $f \in \mathbb{R}[V]$, bounded on S , such that the set $f^{-1}(c) \cap \overline{S}$ is compact for every $0 \neq c \in \mathbb{R}$.*

Note that condition (v) implies that there exists $\varphi: V \rightarrow W$ as in (iv) with $W = \mathbb{A}^1$.

Proof. (i) \Rightarrow (iii): Assume $\text{Quot}(B) = \mathbb{R}(V)$. Choose a finitely generated \mathbb{R} -subalgebra C of B with $\text{Quot}(C) = \mathbb{R}(V)$, and put $W = \text{Spec}(C)$. The morphism $\varphi: V \rightarrow W$ induced by the inclusion $C \subset \mathbb{R}[V]$ satisfies condition (iii).

(iii) \Rightarrow (ii): (iii) implies $\mathbb{R}[W] \hookrightarrow B$, which implies (ii).

(ii) \Rightarrow (i): The field extension $\text{Quot}(B) \subset \mathbb{R}(V)$ is algebraic, by (ii). Therefore, given $f \in \mathbb{R}[V]$, there exists $0 \neq t \in B$ such that tf is integral over B . Since B is integrally closed in $\mathbb{R}[V]$ (Lemma 1.3(e)), this shows $tf \in B$, and hence $f \in \text{Quot}(B)$.

(iii) \Rightarrow (v): For $\varphi: V \rightarrow W$ as in (iii), let D be a proper closed subvariety of W such that the restriction $\varphi^{-1}(W \setminus D) \rightarrow W \setminus D$ of φ is an isomorphism. Choose a non-constant function $g \in \mathbb{R}[W]$ which vanishes on D and on the boundary of $\varphi(S)$ in $W(\mathbb{R})$. Let $0 \neq c \in \mathbb{R}$. Then $g^{-1}(c) \cap \varphi(S) = g^{-1}(c) \cap \overline{\varphi(S)}$ by the choice of g , and this set is compact since $\overline{\varphi(S)}$ is compact. Moreover, $g^{-1}(c) \cap \varphi(S)$ is contained in $(W \setminus D)(\mathbb{R})$, and so the preimage

$$\varphi^{-1}(g^{-1}(c) \cap \varphi(S)) = (g \circ \varphi)^{-1}(c) \cap \varphi^{-1}(\varphi(S))$$

is compact as well. In particular, $(g \circ \varphi)^{-1}(c) \cap \overline{S}$ is compact, and so it suffices to take $f := g \circ \varphi$.

(v) \Rightarrow (iv) is trivial (we can take $W = \mathbb{A}^1$ and $\varphi := f$).

(iv) \Rightarrow (i): Let φ be as in (iv). Given any $f \in \mathbb{R}[V]$, the set $\Omega_\varphi(f)$ is not Zariski dense in W . So Prop. 2.2 shows $f \in \text{Quot}(B)$. \square

Under suitable conditions, we know a priori that the \mathbb{R} -algebra B is finitely generated (see Thm. 5.12 below). In these cases we can formulate Proposition 2.2 and Theorem 2.5 more succinctly:

Corollary 2.6. *Assume that the \mathbb{R} -algebra $B = B_V(S)$ is finitely generated, put $W = \text{Spec}(B)$, and let $\varphi: V \rightarrow W$ be the canonical morphism.*

- (a) $f \in \mathbb{R}[V]$ lies in $\text{Quot}(B)$ if and only if $\Omega_\varphi(f)$ is not Zariski dense in W .
- (b) φ is birational if and only if $\dim(W) = \dim(V)$, if and only if the set Ω_φ is not Zariski dense in $W(\mathbb{R})$. \square

Remark 2.7. Even if $B = B_V(S)$ fails to be finitely generated, we can characterize $\text{trdeg}_{\mathbb{R}}(B)$ as the maximum dimension of an affine \mathbb{R} -variety W for which there exists a dominant morphism $V \rightarrow W$ which is bounded on S .

Examples 2.8.

1. Let $V = \mathbb{A}^2$, and consider the set $S = \{(x, y) : 0 \leq x(x^2 + y^2) \leq 1\}$ in \mathbb{R}^2 . Then $y \notin B(S)$ but $x, xy \in B(S)$, so $y \in \text{Quot}(B)$. In fact, $B = \mathbb{R}[x, xy, xy^2]$. If $\varphi: V \rightarrow W = \text{Spec}(B)$ denotes the canonical map then $\Omega_\varphi = \Omega_\varphi(y)$ consists only of the origin in W .

2. Let $f, g \in \mathbb{R}[x, y]$ be two algebraically independent polynomials, and let $S = \{p \in \mathbb{R}^2 : |f(p)| \leq 1, |f(p)g(p)| \leq 1\}$. Clearly $f, fg \in B(S)$, hence $\text{trdeg } B(S) = 2$. From Theorem 2.5 it follows that there exists a non-constant polynomial $h \in B(S)$ such that all fibres $h^{-1}(c) \cap S$ are compact for $c \neq 0$. Depending on the choice of f and g , it may not be *a priori* clear how to find such h . For a concrete example, take $f = x^2y$ and $g = y^2$. Here, neither f nor fg have any compact fibres, but (for example) $h = xy$ will do.

3. The same example also shows another phenomenon. If $\text{trdeg } B(S) = 2$, there exists a birational morphism $\varphi: \mathbb{A}^2 \rightarrow W$ of affine varieties that is bounded on S . In many instances, the map $\mathbb{A}^2 \rightarrow \mathbb{A}^2, (x, y) \mapsto ((f(x), f(y)g(y)))$ will not have this property, implying in particular that $B(S)$ is strictly larger than $\mathbb{R}[f, fg]$. Finding a birational morphism φ as above can be seen as a first step towards determining $B(S)$. (For f and g as in the previous example, it is easy to see that $B(S) = \mathbb{R}[xy, x^2y, x^2y^3]$, while the map $(x, y) \mapsto (f(x), f(y)g(y))$ has generically degree four.) There does not seem to be any general procedure that will produce such φ . See however Remark 3.7.3.

2.9. Throughout this paper, we shall assume for the most part that varieties are irreducible. Here are a few remarks on the reducible case. Let V be an affine \mathbb{R} -variety with irreducible components V_1, \dots, V_r , let S be a semi-algebraic subset of $V(\mathbb{R})$, and write $S_i := S \cap V_i(\mathbb{R})$. The relation between $B_V(S)$ and the rings $B_{V_i}(S_i)$ ($i = 1, \dots, r$) depends largely on the way the components V_i of V meet. Clearly, the restriction $\mathbb{R}[V] \rightarrow \mathbb{R}[V_i]$ maps $B_V(S)$ to $B_{V_i}(S_i)$ ($i = 1, \dots, r$). But $B_V(S) \rightarrow B_{V_i}(S_i)$ need not be surjective, as the following example shows. Let V be the plane affine curve $x(x^2 + y^2 - 1) = 0$, which is the union of a circle V_1 and a line V_2 , and take $S = V(\mathbb{R})$. Then $B_V(S)$ consists of those polynomials which are constant on the line, and so the restriction map $B_V(S) \rightarrow B_{V_1}(S_1)$ to the circle does not contain $y|_{V_1}$ in its image.

Nevertheless, in the case when V is a curve, the relation between $B_V(S)$ and the $B_{V_i}(S_i)$ is understood fairly well, see [15] for details. Things become considerably more difficult in dimension at least two. For instance, while $B_V(S) = \mathbb{R}$ implies $B_{V_i}(S_i) = \mathbb{R}$ for all i when $\dim(V) = 1$ ([15], Lemma 1.12 (5)), this is false in higher dimensions. For an example, let V be the union of two planes in 3-space, and let S be the union of the first plane $V_1(\mathbb{R})$ with a strip $[0, 1] \times \mathbb{R}$ in the second plane, where the strip is transversal to the line $V_1 \cap V_2$. Then $B_V(S) = \mathbb{R} \neq B_{V_2}(S_2)$.

3. BOUNDED POLYNOMIALS AND COMPLETIONS OF VARIETIES

3.1. Let X be a normal \mathbb{R} -variety. A *prime divisor* on X will be a closed irreducible subset of codimension one in X . By a *divisor* on X we always mean a Weil divisor, that is, an element of the free abelian group generated by the prime divisors on X . Linear equivalence of divisors is denoted \sim . If Z is a prime divisor, the discrete valuation of $\mathbb{R}(X)$ associated with Z will be denoted by v_Z . Thus $v_Z(f)$ is the vanishing order of f along Z , for $f \in \mathbb{R}(X)^*$. Recall that the prime divisor Z is said to be real if its residue field $\mathbb{R}(Z)$ can be ordered (c.f. 1.1).

Given a rational map $f: V \dashrightarrow W$ between irreducible varieties, we denote by $\text{dom}(f)$ the largest open subset of V on which f is defined.

3.2. Let X be an \mathbb{R} -variety. Sometimes it will be convenient to work in the real spectrum X_r of X . When X is affine, $X_r = \text{Sper } \mathbb{R}[X]$ is the space of all orderings of the ring $\mathbb{R}[X]$ (see [3] § 7). When X is not necessarily affine, the topological space X_r is defined by glueing the real spectra $(U_i)_r$ ($i = 1, \dots, r$) of an open affine cover $X = U_1 \cup \dots \cup U_r$, see [17] 0.4 for more details. Thus a point $\alpha \in X_r$ corresponds to a pair (x_α, P_α) where x_α is a (scheme-theoretic) point of X and P_α is an ordering of the residue class field of X in x_α . The support of α is $\text{supp}(\alpha) = \overline{\{x_\alpha\}}$. In particular, $X(\mathbb{R})$ is a topological subspace of X_r in the obvious way. Similarly, when X is irreducible, the space $\text{Sper } \mathbb{R}(X)$ of all orderings of the function field of X is identified with the subspace $\{\alpha \in X_r : \text{supp}(\alpha) = X\}$ of X_r .

The topological space X_r is spectral, and hence there is a well-defined notion of constructible subsets of X_r (see [17] 0.4). For every semi-algebraic subset S of $X(\mathbb{R})$, there exists a unique constructible subset \tilde{S} of X_r such that $S = \tilde{S} \cap X(\mathbb{R})$. If X is affine, then \tilde{S} is the subset of X_r defined by the same system of inequalities as S . It is well-known that \tilde{S} is open (resp. closed) in X_r if and only if the same is true of S in $X(\mathbb{R})$. For two points $\alpha, \beta \in X_r$, we say that α specializes to β , denoted $\alpha \rightsquigarrow \beta$, if β is contained in the closure of α .

Recall that a valuation v of a field K is called *compatible* with an ordering \leq of K if $0 < b \leq a$, for $a, b \in K^*$, implies $v(b) \geq v(a)$. The usefulness of the real spectrum in our context comes from the following lemma:

Lemma and Definition 3.3. *Let X be a normal \mathbb{R} -variety, let Z be a prime divisor on X , and let S be a semi-algebraic subset of $X(\mathbb{R})$. The following conditions are equivalent:*

- (i) $Z(\mathbb{R}) \cap \overline{(S \cap U(\mathbb{R}))}$ is Zariski dense in Z , where $U = X \setminus Z$;
- (ii) there is a specialization $\alpha \rightsquigarrow \beta$ in X_r with $\alpha \in \tilde{S}$, $\text{supp}(\beta) = Z$ and $\alpha \neq \beta$;
- (iii) the discrete valuation v_Z of $\mathbb{R}(X)$ is compatible with some ordering in $\tilde{S} \cap \text{Sper } \mathbb{R}(X)$.

If these conditions hold we say that Z and S are compatible.

Note that every prime divisor which is compatible with S has a real residue field.

Proof. Let $T := Z(\mathbb{R}) \cap \overline{(S \cap U(\mathbb{R}))}$, a closed semi-algebraic subset of $Z(\mathbb{R})$. The set T is Zariski dense in Z if and only if \tilde{T} contains a point with support Z . The latter condition is equivalent to (ii), and (iii) is merely a reformulation of (ii). \square

The following lemma is obvious (see [18] Lemma 0.2) and will be used frequently:

Lemma 3.4. *Let V be a connected normal \mathbb{R} -variety, let $S \subset V(\mathbb{R})$ be a semi-algebraic set, and let Z be a prime divisor in V which is compatible with S . Let $f_1, \dots, f_r \in \mathbb{R}(V)$ satisfy $f_i \geq 0$ on $S \cap \text{dom}(f)$. Then*

$$v_Z\left(\sum_i f_i\right) = \min_i v_Z(f_i). \quad \square$$

Lemma 3.5. *Let V be a normal \mathbb{R} -variety, let $S \subset V(\mathbb{R})$ be a semi-algebraic set, and let Z be a prime divisor in V which is compatible with S . If a rational function $f \in \mathbb{R}(V)^*$ has a pole along Z , then f is unbounded on $S \cap \text{dom}(f)$.*

Proof. The compatibility of Z with S means that there exists $\alpha \in \tilde{S} \cap \text{Sper } \mathbb{R}(V)$ which makes the discrete valuation ring $\mathcal{O}_{V,Z}$ convex in $\mathbb{R}(V)$. Assume that f is bounded on $S' := S \cap \text{dom}(f)$. Since \tilde{S}' has the same trace in $\text{Sper } \mathbb{R}(V)$ as \tilde{S} , we have $\alpha \in \tilde{S}'$ as well. On the other hand, f bounded on S' implies that f lies in the α -convex hull of \mathbb{R} in $\mathbb{R}(V)$. In particular, $f \in \mathcal{O}_{V,Z}$, which means that f does not have a pole along Z . \square

Definition 3.6. Let V be an irreducible \mathbb{R} -variety and let S be a semi-algebraic subset of $V(\mathbb{R})$. By a *completion* of V we mean an open dense embedding $V \hookrightarrow X$ into a complete \mathbb{R} -variety. The completion X will be said to be *compatible with S* (or *S -compatible*) if for every irreducible component Z of $X \setminus V$ the following conditions hold:

- (1) The local ring $\mathcal{O}_{X,Z}$ is a discrete valuation ring;
- (2) the set $Z(\mathbb{R}) \cap \bar{S}$ is either empty or Zariski dense in Z .

(Here, of course, \bar{S} denotes the closure of S in $X(\mathbb{R})$.)

Note that (1) is saying that Z has codimension one in X and is not contained in the singular locus of X . Condition (2) says (for normal X) that every irreducible component of $X \setminus V$ is either compatible with S or disjoint from \bar{S} (c.f. Definition 3.3).

Examples 3.7.

1. Given a semi-algebraic subset S of \mathbb{R}^n , the natural completion $\mathbb{A}^n \subset \mathbb{P}^n$ of affine n -space is compatible with S if and only if S contains an open cone in \mathbb{R}^n (not necessarily centered at the origin).

2. The case of curves is simple: Given a (possibly singular) irreducible curve C , there is a unique projective completion $C \hookrightarrow X$ for which $X_{\text{sing}} \subset C_{\text{sing}}$ (see [19] 4.6). The completion X is compatible with any semi-algebraic subset S of $C(\mathbb{R})$. The points of $X \setminus C$ are called the *points of C at infinity*.

3. There are interesting classes of semi-algebraic sets S (in \mathbb{R}^n , say) for which S -compatible completions (of $V = \mathbb{A}^n$, in this case) can be constructed as toric varieties. For example, when S is defined by (finitely many) binomial inequalities $ax^\alpha < bx^\beta$, this always is the case. For such S , the ring $B(S)$ can be identified

explicitly in terms of the defining inequalities, and $B(S)$ is always finitely generated as an \mathbb{R} -algebra. We plan to expand on this remark in a future publication.

Our interest in compatible completions arises from the next result. It shows that such a completion calculates the ring of bounded polynomials on a semi-algebraic set:

Theorem 3.8. *Let V be an affine normal \mathbb{R} -variety, let $S \subset V(\mathbb{R})$ be a semi-algebraic subset, and assume that the completion $V \hookrightarrow X$ of V is compatible with S . Let Y denote the union of those irreducible components Z of $X \setminus V$ for which $\overline{S} \cap Z(\mathbb{R}) = \emptyset$, and put $U = X \setminus Y$. Then the inclusion $V \subset U$ induces a ring isomorphism*

$$\mathcal{O}_X(U) \xrightarrow{\sim} B_V(S).$$

Proof. Again, \overline{S} denotes the closure of S in $X(\mathbb{R})$. Since \overline{S} is compact and contained in $U(\mathbb{R})$, every element of $\mathcal{O}_X(U)$ is bounded on S . So the image of the restriction map $\mathcal{O}_X(U) \hookrightarrow \mathbb{R}[V]$ is contained in $B_V(S)$. It remains to show that every $f \in B_V(S)$, considered as a rational function on U , is regular on U . Since U is normal, it suffices that $v_Z(f) \geq 0$ for every irreducible component Z of $U \setminus V$. By the construction of U , and since X is compatible with S , the intersection $\overline{S} \cap Z(\mathbb{R})$ is Zariski dense in Z , which means that the divisor Z is compatible with the set S (see 3.3). Since f is regular on V and bounded on S , Lemma 3.5 implies $v_Z(f) \geq 0$. \square

3.9. We illustrate the possible use of Thm. 3.8 by two examples. We use homogeneous coordinates $(x_0 : x_1 : x_2)$ on \mathbb{P}^2 , and we identify $(x, y) \in \mathbb{A}^2$ with $(1 : x : y) \in \mathbb{P}^2$. Let $L = \{x_0 = 0\}$ be the line at infinity in \mathbb{P}^2 .

Consider the strip $S = \{(x, y) : |x| \leq 1\}$ in \mathbb{R}^2 . In $\mathbb{P}^2(\mathbb{R})$ we have $\overline{S} \cap L(\mathbb{R}) = \{P\}$ where $P = (0 : 0 : 1)$. To improve this we consider the blowing-up $\pi_1 : X_1 \rightarrow \mathbb{P}^2$ at P . Then $\mathbb{A}^2 \hookrightarrow X_1$ is a completion of \mathbb{A}^2 for which $X_1 \setminus \mathbb{A}^2$ has two irreducible components, namely L' (the strict transform of L) and $E_1 = \pi_1^{-1}(P)$. In $X_1(\mathbb{R})$ we have $\overline{S} \cap L'(\mathbb{R}) = \emptyset$, and $\overline{S} \cap E_1(\mathbb{R})$ is Zariski dense in E_1 . Therefore the completion X_1 is compatible with S , and by 3.8 we have $B(S) = \mathcal{O}(U)$ for $U := X_1 \setminus L'$. Clearly $\mathcal{O}(U) = \{f \in \mathbb{R}[x, y] : v_{E_1}(f) \geq 0\}$. Calculating the valuation v_{E_1} we find

$$v_{E_1} \left(\sum_{i,j} a_{ij} x^i y^j \right) = \min\{-j : a_{ij} \neq 0\}.$$

So we get $B(S) = \mathcal{O}(U) = \mathbb{R}[x]$.

3.10. For another example let now $T = \{(x, y) \in \mathbb{R}^2 : |x| \leq 1, |xy| \leq 1\}$. As before we use the blowing-up $\pi_1 : X_1 \rightarrow \mathbb{P}^2$ of the plane in $P = (0 : 0 : 1)$. Again we have $\overline{T} \cap L'(\mathbb{R}) = \emptyset$ in $X_1(\mathbb{R})$, but this time $\overline{T} \cap E_1(\mathbb{R}) = \{P_1\}$ is a singleton. Therefore we blow up X_1 in P_1 to get $\pi_2 : X_2 \rightarrow X_1$, with exceptional fibre $E_2 = \pi_2^{-1}(P_1)$. Then in $X_2(\mathbb{R})$ we find $\overline{T} \cap E_1'(\mathbb{R}) = \emptyset$, and $\overline{T} \cap E_2(\mathbb{R})$ is Zariski dense in E_2 . So X_2 is a completion of \mathbb{A}^2 which is compatible with T , and $B(T) = \mathcal{O}(W)$ for $W := X_2 \setminus (L'' \cup E_1')$ according to Thm. 3.8. We find $\mathcal{O}(W) = \{f \in \mathbb{R}[x, y] : v_{E_2}(f) \geq 0\}$ and

$$v_{E_2} \left(\sum_{i,j} a_{ij} x^i y^j \right) = \min\{i - j : a_{ij} \neq 0\},$$

which shows $B(T) = \mathcal{O}(W) = \mathbb{R}[x, xy]$.

In general, if we want to apply Theorem 3.8 to calculate $B_V(S)$, we can start with some (normal) completion $V \hookrightarrow X_0$ of V . By making suitable iterated blowing-ups with centers over $X_0 \setminus V$, we try to “straighten out” the set S at infinity more and more. When V is a surface and the set S is sufficiently regular at infinity, this procedure will always lead, after finitely many steps, to a completion of V which is compatible with S , see Theorem 4.5 below.

4. EXISTENCE OF COMPATIBLE COMPLETIONS

In view of Theorem 3.8, we are now discussing the existence question for S -compatible completions. We start with the case $S = V(\mathbb{R})$.

Theorem 4.1. *Every connected non-singular \mathbb{R} -variety V has a completion X which is compatible with the set $S = V(\mathbb{R})$, and such that the irreducible components of $X \setminus V$ are non-singular. If V is quasi-projective then X can be chosen to be projective.*

Proof. Start with any open dense embedding $V \hookrightarrow X$ into a complete \mathbb{R} -variety X (which we can take projective if V is quasi-projective). The singularities of X are contained in $X \setminus V$, and by resolving them we get X non-singular. Some irreducible components of $X \setminus V$ may have codimension ≥ 2 . This can be remedied by blowing up X in these centers. So we can assume that every irreducible component of $X \setminus V$ has codimension one in X . Finally, by embedded resolution of singularities, we can achieve that the irreducible components of $X \setminus V$ are non-singular.

We claim that the completion $V \subset X$ is compatible with $V(\mathbb{R})$. To see this, fix an irreducible component Z of $X \setminus V$ for which $Z(\mathbb{R}) \neq \emptyset$. Since Z is non-singular, the function field $\mathbb{R}(Z)$ is real. By the Baer-Krull theorem, $\mathbb{R}(X)$ has an ordering α which is compatible with the discrete valuation v_Z . In particular, Z is compatible with the set $V(\mathbb{R})$ (Definition 3.3). \square

We now turn to compatible completions for more general semi-algebraic subsets S of $V(\mathbb{R})$. We shall denote the interior of a set $M \subset V(\mathbb{R})$ by $\text{int}(M)$.

Definition 4.2. Let V be an irreducible \mathbb{R} -variety, let $S \subset V(\mathbb{R})$ be a semi-algebraic set.

- (a) The set S is said to be *regular* if $S \subset \overline{\text{int}(S \cap V_{\text{reg}}(\mathbb{R}))}$.
- (b) S is called *regular at infinity* if $S = S_0 \cup S_1$ where S_0, S_1 are semi-algebraic sets with \bar{S}_0 compact and S_1 regular.
- (c) S is called *Zariski dense at infinity* if $S \setminus (K \cap S)$ is Zariski dense in V for every compact subset K of $V(\mathbb{R})$.

Remark 4.3. If S is regular at infinity and \bar{S} is not compact, then S is Zariski dense at infinity. Indeed, otherwise there would be a proper Zariski closed subset Z of V and a compact set $K \subset V(\mathbb{R})$ with $S \subset K \cup Z(\mathbb{R})$. This would imply $\text{int}(S \cap V_{\text{reg}}(\mathbb{R})) \subset K$, and by regularity at infinity we would get \bar{S} compact, a contradiction.

We will need the following version of embedded resolution of singularities on a surface. By a curve on X we mean a reduced effective divisor on X .

Theorem 4.4. *Let k be an infinite field, X a normal quasi-projective surface over k , C a curve in X and T a finite set of closed points in $C_{\text{sing}} \cap X_{\text{reg}}$. Then there exists a birational morphism $\varphi: \tilde{X} \rightarrow X$ of k -varieties with the following properties:*

- (1) φ induces an isomorphism $\tilde{X} \setminus \varphi^{-1}(T) \xrightarrow{\sim} X \setminus T$;
- (2) \tilde{X} is quasi-projective and $\tilde{X}_{\text{sing}} = \varphi^{-1}(X_{\text{sing}})$ (in particular, \tilde{X} is normal);
- (3) in all points of $\varphi^{-1}(T)$, the divisor $\varphi^{-1}(C)$ on \tilde{X} has normal crossings and non-singular components.

Proof. If X is non-singular and $T = C_{\text{sing}}$, our statement becomes the usual one for embedded resolution of curves in surfaces (see for example [6] Thm. V.3.7, or [4] Sect. 3.5 for the case of an arbitrary infinite base field). This implies the above version, since T is contained in X_{reg} . In more detail, let $S = X_{\text{sing}} \cup (C_{\text{sing}} \setminus T)$ and put $X_0 = X \setminus S$, a non-singular quasi-projective surface. Applying usual embedded resolution of singularities to the divisor $C \cap X_0$ on X_0 and glueing it with the identity of $X \setminus T$ yields the above version. \square

The following result proves the existence of compatible completions for surfaces, when the semi-algebraic set is regular at infinity (Definition 4.2):

Theorem 4.5. *Let V be a connected normal quasi-projective surface over \mathbb{R} , and let S be a semi-algebraic subset of $V(\mathbb{R})$ that is regular at infinity. Then V has a projective completion which is compatible with S . If V is non-singular then the completion can be chosen to be non-singular as well.*

Proof. We may assume that the function field $\mathbb{R}(V)$ is real. For otherwise, $V(\mathbb{R})$ consists of finitely many singular points of V . In that case, any normal completion of V will be compatible with any subset of $V(\mathbb{R})$.

We can also assume that S is closed in $V(\mathbb{R})$. Start with any completion $V \hookrightarrow X_1$ of V . Since V is a normal surface, the singular set V_{sing} is finite. By resolving singularities of $X_1 \setminus V_{\text{sing}}$ and gluing the resulting surface to V we can assume $X_{1,\text{sing}} \subset V$. Let \bar{S} be the closure of S in $X_1(\mathbb{R})$, let $\partial\bar{S}$ be its boundary in $X_1(\mathbb{R})$, and denote by C_1 the Zariski closure of $\partial\bar{S}$ in X_1 , a curve on X_1 . Further write $D_1 = X_1 \setminus V$. Now apply embedded resolution, as stated in Theorem 4.4, to the surface X_1 , the curve $C_1 \cup D_1$ on X and the set $T := (C_1 \cup D_1)_{\text{sing}} \cap D_1$. We obtain a birational morphism $\pi: X \rightarrow X_1$ whose restriction $\pi^{-1}(V) \rightarrow V$ is an isomorphism. Via π we can regard X as a completion of V . Let $D = X \setminus V = \pi^{-1}(D_1)$ and $C = \pi^{-1}(C_1)$. The completion $V \hookrightarrow X$ has the following properties:

- (1) X is projective and normal (non-singular if V is non-singular);
- (2) the irreducible components of the divisor $C \cup D$ are non-singular in all points of D , and if two components of $C \cup D$ meet in a point of D , then that point is a normal crossing singularity.

We show that $V \hookrightarrow X$ is compatible with S . Let Z be an irreducible component of $D = X \setminus V$ for which the set $\bar{S} \cap Z(\mathbb{R})$ is non-empty. We have to show that this set is Zariski dense in Z . Let $x \in \bar{S} \cap Z(\mathbb{R})$, and let $u \in \mathcal{O}_{X,x}$ be a local equation for Z . There can be at most one more irreducible component $Z' \neq Z$ of $C \cup D$ which passes through x , by property (2). If such Z' exists, let $v \in \mathcal{O}_{X,x}$ be a local equation for Z' . Then u, v is a regular system of parameters for $\mathcal{O}_{X,x}$. If no such Z' exist we put $Z' = Z$ and let $v \in \mathcal{O}_{X,x}$ be any element for which u, v is a regular system of parameters for $\mathcal{O}_{X,x}$.

Let U be an open neighbourhood of x in $X(\mathbb{R})$ such that $U \cap (C \cup D)(\mathbb{R}) \subset (Z \cup Z')(\mathbb{R})$. By shrinking U we can assume that u and v are regular functions on U . Shrinking U further if necessary, there are precisely four connected components of $\{y \in U : (uv)(y) \neq 0\}$. Since S is regular at infinity, it follows that \bar{S} contains

at least one of these four components. In particular, $\overline{S} \cap Z(\mathbb{R})$ contains a non-empty open subset of $Z(\mathbb{R})$ and is therefore Zariski dense in Z . This completes the proof. \square

Remarks 4.6.

1. We do not know whether Theorem 4.5 extends to $\dim(V) > 2$. See however Theorem 4.1.

2. In Theorem 4.5, the regularity of S at infinity is not necessary in order that a compatible completion exists. This is demonstrated by simple examples like the following: Let $V = \mathbb{A}^2$ and consider the set

$$S = \{(x, y) \in \mathbb{R}^2 : xy^2 \geq 0\}.$$

S is the union of the right half plane with the x -axis. Clearly \mathbb{P}^2 is a completion of \mathbb{A}^2 which is compatible with S , even though S is not regular at infinity. On the other hand, when $S \subset V(\mathbb{R})$ is unbounded but contained in the union of a compact set and a proper subvariety of V (so S fails to be Zariski dense at infinity), there cannot be any completion of V compatible with S (see Prop. 5.7.)

3. If C is a possibly singular (irreducible) affine curve, it is still true that C has a unique projective completion X which is non-singular in the added points (c.f. also [19] Lemma 4.5). The completion X is compatible with every semi-algebraic subset S of $C(\mathbb{R})$, and Theorem 3.8 holds in this case as well.

4. Let X be a non-singular connected projective surface over \mathbb{R} , and let D be a curve on X with irreducible components Z_1, \dots, Z_r . Then it is sometimes possible to calculate the transcendence degree of the ring $B = \mathcal{O}_X(X \setminus D)$ from the intersection matrix $M_D = (Z_i \cdot Z_j)_{i,j=1,\dots,r}$ of the divisor D :

- (1) If M_D is negative definite, then $\text{trdeg}_{\mathbb{R}}(B) = 0$, hence $B = \mathbb{R}$.
- (2) If M_D has a positive eigenvalue, then $\text{trdeg}_{\mathbb{R}}(B) = 2$.

If M_D is negative semidefinite and singular, there is no general statement about the transcendence degree of B . (See [7] 8.3 for proofs.)

For instance, consider $T = \{(x, y) \in \mathbb{R}^2 : |x| \leq 1, |xy| \leq 1\}$ as in Example 3.10. There we found $B(T) = \mathcal{O}(X_2 \setminus D)$ with $D = L'' \cup E'_1$ (using the notation from 3.10). The intersection matrix of D is

$$M_D = \begin{pmatrix} 0 & 1 \\ 1 & -2 \end{pmatrix},$$

which has a positive eigenvalue. Thus we conclude $\text{trdeg } B(T) = 2$ without actually calculating the ring $B(T)$ as in 3.10. In this example, the transcendence degree can also be read off using Thm. 2.5, since the birational map $\mathbb{A}^2 \rightarrow \mathbb{A}^2$, $(x, y) \mapsto (x, xy)$ is bounded on S .

Let V be an affine normal \mathbb{R} -variety and let $S \subset V(\mathbb{R})$ be a semi-algebraic subset. We have seen that the ring $B_V(S)$ is isomorphic to a ring $\mathcal{O}_U(U)$ for some quasi-projective \mathbb{R} -variety U , provided that V has a completion which is compatible with S . We are now going to prove a converse (Thm. 4.11 below). It roughly states, for every quasiprojective U which is real, that the ring $\mathcal{O}_U(U)$ can be realized as $B_V(S)$ for some open affine subset V of U and some semi-algebraic subset S of $V(\mathbb{R})$.

Lemma 4.7. *Let V be an irreducible affine \mathbb{R} -variety, and let X be a projective completion of V which is normal. Let Y_1, \dots, Y_r be irreducible components of $X \setminus V$ which are real, and let Z_1, \dots, Z_s be the remaining components (real or*

not) of $X \setminus V$. There exists a regular and basic closed semi-algebraic subset S of $V(\mathbb{R})$ that is compatible with Y_1, \dots, Y_r and whose closure in $X(\mathbb{R})$ is disjoint from $Z_1(\mathbb{R}), \dots, Z_s(\mathbb{R})$. In particular, the completion X of V is compatible with S .

Proof. Fix an index $i \in \{1, \dots, r\}$. Since X is normal and Y_i is real, we find a point $p_i \in Y_i(\mathbb{R})$ which is non-singular on X and on $X \setminus V$. By this last fact, p_i has a generalisation β_i in $\text{Sper } \mathbb{R}(Y_i)$. Since X is normal, β_i has a generalization α_i in $\text{Sper } \mathbb{R}(X)$. Every semi-algebraic subset S of $X(\mathbb{R})$ with $\alpha_i \in \tilde{S}$ is compatible with Y_i . There is an open affine subset U of X such that $U(\mathbb{R}) = X(\mathbb{R})$. Let x_1, \dots, x_n be a system of generators for the \mathbb{R} -algebra $\mathbb{R}[U]$. If $c_i \in \mathbb{R}$ is positive and sufficiently small, the closed semi-algebraic set

$$S_i := \left\{ p \in V(\mathbb{R}) : \sum_{k=1}^n (x_k(p) - x_k(p_i))^2 \leq c_i \right\}$$

will be contained in $V_{\text{reg}}(\mathbb{R})$ and regular, and moreover S_i will be compatible with Y_i and disjoint from $Z_1(\mathbb{R}) \cup \dots \cup Z_s(\mathbb{R})$. This being done for $i = 1, \dots, r$, we may further assume that S_1, \dots, S_r are pairwise disjoint, by making c_1, \dots, c_r even smaller if necessary.

We claim that $S := S_1 \cup \dots \cup S_r$ is a basic closed semi-algebraic subset of $V(\mathbb{R})$, which will complete the proof. For $i = 1, \dots, r$ put

$$g_i := c_i - \sum_{k=1}^n (x_k - x_k(p_i))^2 \in \mathbb{R}[U],$$

and let V_0 be the maximal Zariski open subset of V for which $g_i|_{V_0} \in \mathcal{O}_V(V_0)$ for $i = 1, \dots, r$. Since V is normal, the closed subset $Z = V \setminus V_0$ has pure codimension one in V . Clearly $Z(\mathbb{R}) = \emptyset$ since $U(\mathbb{R}) = X(\mathbb{R})$ and the g_i are regular on U . Choose $h_1, \dots, h_k \in \mathbb{R}[V]$ which generate the vanishing ideal of Z in $\mathbb{R}[V]$, and put $h := h_1^2 + \dots + h_k^2$. Then h has no real zeros on V , and since V is normal, there exists $N \geq 1$ such that $f_i := h^N g_i \in \mathbb{R}[V]$ for $i = 1, \dots, r$. Thus $S_i = \{p \in V(\mathbb{R}) : f_i(p) \geq 0\}$ for every i , and hence

$$S_1 \cup \dots \cup S_r = \{p \in V(\mathbb{R}) : (-1)^{r+1} f_1(p) \cdots f_r(p) \geq 0\},$$

since the S_i are pairwise disjoint. \square

An effective Weil divisor D on a normal \mathbb{R} -variety X will be called *totally real* if every irreducible component of D is a real variety. By $\text{Cl}(X)$ we denote the class group of Weil divisors modulo linear equivalence on X . We will use the following theorem:

Theorem 4.8 (Roggero [16]). *Let V be a normal affine real \mathbb{R} -variety of dimension at least 2. Then every divisor on V is linearly equivalent to a totally real effective divisor on V .* \square

Corollary 4.9. *Let X be a real normal projective \mathbb{R} -variety of dimension at least 2, and let H be an ample effective divisor on X . For every divisor D on X there exists a totally real effective divisor D_0 on X such that $D \sim D_0 + H_0$, where H_0 is a linear combination of irreducible components of H .*

Proof. Let $V = X \setminus H$. The kernel of the restriction map $\text{Cl}(X) \rightarrow \text{Cl}(V)$ is generated by (the class of) H . Thus it suffices to apply Roggero's theorem to the affine variety V . \square

Lemma 4.10. *Let U be a connected normal quasi-projective \mathbb{R} -variety which is real. There exist real prime divisors Y_1, \dots, Y_r on U such that the variety $U \setminus (Y_1 \cup \dots \cup Y_r)$ is affine.*

Proof. If U is a curve, we can take $r = 1$ and Y_1 any real point on U . So we may assume $\dim(U) \geq 2$. Assume furthermore that U is projective. Fix a non-singular point $p \in U(\mathbb{R})$ and an ample divisor H on U , and let $L \subset |H|$ consist of those $D \in |H|$ that pass through p . The linear system L has no fixed components, so by Bertini's theorem (see for example Jouanolou [8], Cor. 6.11), the set of all $Z \in L$ for which Z is irreducible and satisfies $Z_{\text{sing}} \subset U_{\text{sing}}$ has non-empty interior inside L . In particular, it is non-empty. Any such Z is real since it contains p as a regular real point. Moreover $U \setminus Z$ is affine since Z is ample.

When U is only quasi-projective, fix an open dense embedding $U \hookrightarrow X$ with X normal and projective and such that $X \setminus U$ has pure codimension one in X (the latter can be achieved by blowing up if necessary). Let E be the divisor $X \setminus U$. By the projective case above we find an ample prime divisor H on X that is totally real. By Cor. 4.9, there exist $n \in \mathbb{Z}$ and a totally real effective divisor D on X such that $-E \sim D + nH$. For $m > \max\{0, n\}$ the divisor $E + D + mH$ is effective and ample. It follows that the irreducible components Y_1, \dots, Y_r of $(D + mH) \cap U$ are real and $U \setminus (Y_1 \cup \dots \cup Y_r)$ is affine, as desired. \square

Theorem 4.11. *Let U be a connected normal quasi-projective \mathbb{R} -variety which is real. There exists an open affine subset V of U and a regular and basic closed subset S of $V(\mathbb{R})$ such that $\mathcal{O}_U(U) \xrightarrow{\sim} B_V(S)$.*

Proof. Let $U \hookrightarrow X$ be a normal projective completion of U , and let Z_1, \dots, Z_s be the irreducible components of $X \setminus U$. By Lemma 4.10 there exist further prime divisors Y_1, \dots, Y_r on X which are real and such that the open subset $V := X \setminus (\bigcup_i Y_i \cup \bigcup_j Z_j)$ of U is affine. By Lemma 4.7 we find a regular and basic closed semi-algebraic set S in $V(\mathbb{R})$ which is compatible with Y_1, \dots, Y_r and which satisfies $S \cap Z_j(\mathbb{R}) = \emptyset$ for $j = 1, \dots, s$. In particular, the completion X of V is compatible with S . Therefore $\mathcal{O}_U(U) \xrightarrow{\sim} B_V(S)$ by Theorem 3.8. \square

Under certain conditions we know that the \mathbb{R} -algebra $B_V(S)$ is finitely generated, see e.g. Thm. 5.12 below. Writing $W := \text{Spec } B_V(S)$ for the affine \mathbb{R} -variety defined by this ring, the canonical morphism $\varphi_S: V \rightarrow W$ is universal for morphisms that are bounded on S from V to affine varieties, i.e., every morphism $V \rightarrow W'$ into an affine \mathbb{R} -variety W' which is bounded on S factors uniquely through φ_S . Note that the variety W is normal, provided V is normal (Lemma 1.3(e)). We are wondering how to characterize all morphisms $V \rightarrow W$ of normal affine \mathbb{R} -varieties which are of the form $\varphi_S: V \rightarrow \text{Spec } B_V(S)$ for some semi-algebraic subset S of $V(\mathbb{R})$ (with $B_V(S)$ finitely generated).

Although we do not know a general answer to this question, we can present the following sufficient condition for birational morphisms:

Proposition 4.12. *Let $\varphi: V \rightarrow W$ be a birational morphism of connected normal real affine \mathbb{R} -varieties. Assume there is a totally real effective divisor Z on W such that φ restricts to an isomorphism $\varphi^{-1}(W \setminus Z) \xrightarrow{\sim} W \setminus Z$. Then there exists a semi-algebraic set $S \subset V(\mathbb{R})$ such that $\varphi^*\mathbb{R}[W] = B_V(S)$.*

For the proof we need the following lemma:

Lemma 4.13. *Let V be an affine normal \mathbb{R} -variety, and let Z be a given effective divisor on V which is totally real. There exists a compact semi-algebraic set S in $V(\mathbb{R})$ which is compatible with every irreducible component of Z , and such that $S \setminus (S \cap Z(\mathbb{R}))$ is dense in S .*

Proof. Let Z_1, \dots, Z_r be the irreducible components of Z , and fix $i \in \{1, \dots, r\}$. Since Z_i is real and not contained in V_{sing} , we find a point $p_i \in Z_i(\mathbb{R})$ which is nonsingular on Z_i and on V . So p_i has a generalization β_i in $\text{Sper } \mathbb{R}(Z_i)$, and in turn, β_i has a generalization α_i in $\text{Sper } \mathbb{R}(V)$. Every semi-algebraic subset S of $V(\mathbb{R})$ with $\alpha_i \in \tilde{S}$ is compatible with Z_i . If x_1, \dots, x_n is a system of generators of the \mathbb{R} -algebra $\mathbb{R}[V]$, we may therefore define S to be the closure of the set $\{p \in V(\mathbb{R}) : p \notin Z(\mathbb{R}), x_1(p)^2 + \dots + x_n(p)^2 < c\}$, where $c \in \mathbb{R}$ is positive and sufficiently large. \square

Proof of Prop. 4.12. By Lemma 4.13 there exists a compact basic closed semi-algebraic subset T of $W(\mathbb{R})$ which is compatible with every irreducible component of the divisor Z , and such that $T \setminus (T \cap Z(\mathbb{R}))$ is dense in T . Let $S := \overline{\varphi^{-1}(T)} \subset V(\mathbb{R})$. Since $\varphi(S)$ contains $T \setminus (T \cap Z(\mathbb{R}))$ by the hypothesis, we have $\varphi(S) = T$. By compactness of T it is clear that $\varphi^*(\mathbb{R}[W]) \subset B_V(S)$. We claim that this inclusion is an equality. Let $f \in \mathbb{R}[V]$ with $f \notin \varphi^*(\mathbb{R}[W])$. Considering f as a rational function on W , it follows that f has a pole along one of the irreducible components of Z , since W is normal. Since T is compatible with that component, it follows from Lemma 3.5 that f is unbounded on $T \cap \text{dom}_W(f)$. In particular, f is unbounded on $T \setminus (T \cap Z(\mathbb{R}))$, and so f is unbounded on S . \square

5. FINITE GENERATION OF THE RING OF BOUNDED POLYNOMIALS

We first recall that $B_C(S)$ is always finitely generated when C is a curve:

Proposition 5.1. *Let C be an irreducible affine curve over \mathbb{R} , possibly singular, and let $S \subset C(\mathbb{R})$ be a semi-algebraic subset. Assume that C is real.*

- (a) $B_C(S)$ is finitely generated as an \mathbb{R} -algebra.
- (b) If every point of C at infinity is real and lies in the closure of S , then $B_C(S) = \mathbb{R}$.
- (c) Otherwise $B_C(S)$ has transcendence degree one over \mathbb{R} .

Proof. Let $C \hookrightarrow X$ be the completion of C for which $X_{\text{sing}} \subset C_{\text{sing}}$ (3.7.2), and let $C' \subset X$ be the open subset which is the union of C and all \mathbb{R} -points of $X \setminus C$ which lie in the closure of S . Then $\mathcal{O}_X(C') \xrightarrow{\sim} B_C(S)$. Indeed, even though Thm. 3.8 is not directly applicable here when C is singular, an inspection of the proof shows that it applies nevertheless. When $C' \neq X$ then the curve C' is affine, and so $B_C(S)$ is a finitely generated \mathbb{R} -algebra of transcendence degree one. When $C' = X$ (which means that the hypothesis of (b) is satisfied) then $B_C(S) = \mathbb{R}$. \square

Before proceeding to our main result on finite generation, we study some consequences for the ring $B_V(S)$ that arise from the existence of an S -compatible completion.

Lemma and Definition 5.2. *Let V be an irreducible \mathbb{R} -variety, and let S be a semi-algebraic subset of $V(\mathbb{R})$. There is a unique smallest Zariski closed subset Z of V such that $S \subset Z(\mathbb{R}) \cup K$ for some compact subset K of $V(\mathbb{R})$. We shall call Z the Zariski closure of S at infinity.*

Proof. If Zariski closed subsets Z_1, Z_2 of V and compact subsets K_1, K_2 of $V(\mathbb{R})$ are given with $S \subset Z_i(\mathbb{R}) \cup K_i$ ($i = 1, 2$), then S is contained in $(Z_1(\mathbb{R}) \cap Z_2(\mathbb{R})) \cup (K_1 \cup K_2)$. This implies the assertion. \square

Clearly, S is Zariski dense at infinity (Definition 4.2) if and only if its Zariski closure at infinity is V .

Proposition 5.3. *Let V be an irreducible affine \mathbb{R} -variety and $S \subset V(\mathbb{R})$ a semi-algebraic set. Let Z be the Zariski closure of S at infinity, and let I_Z be the full vanishing ideal of Z in $\mathbb{R}[V]$. Then I_Z is equal to the conductor of $B_V(S)$ in $\mathbb{R}[V]$, that is, I_Z is the largest ideal of $\mathbb{R}[V]$ which is contained in $B_V(S)$.*

Proof. Since $S \subset Z(\mathbb{R}) \cup K$ for some compact set K , it is clear that $I_Z \subset B_V(S)$. Conversely let $b \in B_V(S)$ with $b \notin I_Z$. The subset $\{x \in S : b(x) \neq 0\}$ of $V(\mathbb{R})$ is unbounded. For if its closure K were compact, we would have $S \subset K \cup \{b = 0\}$, contradicting the definition of Z . Hence there exists a semi-algebraic curve $\gamma : [0, \infty[\rightarrow S$, $t \mapsto \gamma_t$ such that $b(\gamma_t) \neq 0$ for all $t \geq 0$ and such that $\Gamma := \{\gamma_t : t \geq 0\}$ is unbounded in $V(\mathbb{R})$. This last condition means that there exists $f \in \mathbb{R}[V]$ for which $f(\gamma_t)$, $t \geq 0$, is unbounded. Since $b \neq 0$ on Γ , it follows from Lemma 5.4 below that the product $f^n b$ is unbounded on Γ for sufficiently large $n \geq 1$. So $b f^n \notin B_V(S)$. \square

The following well-known fact was used in the last proof:

Lemma 5.4. *Let $f, g : [0, \infty[\rightarrow \mathbb{R}$ be two continuous semi-algebraic functions such that f is unbounded and $g > 0$ everywhere. Then $f^n g$ is unbounded for sufficiently large $n \geq 1$.* \square

Lemma 5.5. *Let $B \subset A$ be a ring extension, and let J be an ideal of A which is contained in B . If J contains an element which is not a zero divisor of A , and if J is finitely generated as an ideal of B , then the extension $B \subset A$ is integral.*

Proof. This is a special case of [5, Cor. 4.6]. \square

Lemma 5.6. *Let Y be a connected normal \mathbb{R} -variety, and let $V \subset Y$ be an affine open subset. If $V \neq Y$ then $\mathcal{O}_Y(Y)$ contains no non-zero ideal of $\mathbb{R}[V]$.*

Proof. The inclusion $\mathcal{O}(Y) \subset \mathbb{R}[V]$ is proper. Indeed, otherwise there would be a morphism $Y \rightarrow V$ which is the identity on V , which is absurd if $V \neq Y$. Assume $b \in \mathcal{O}(Y)$ is such that $b f \in \mathcal{O}(Y)$ for every $f \in \mathbb{R}[V]$. Choose $f \in \mathbb{R}[V]$ with $f \notin \mathcal{O}(Y)$. Since Y is normal this means that there exists a prime divisor Z of Y with $v_Z(f) \leq -1$. Unless $b = 0$ we therefore get $v_Z(b f^n) \leq -1$ for large $n > 0$, and so $b f^n \notin \mathcal{O}(Y)$. This proves the assertion. \square

Proposition 5.7. *Let V be a connected normal affine \mathbb{R} -variety, and let S be an unbounded semi-algebraic subset of $V(\mathbb{R})$. Consider the following conditions:*

- (i) V has a completion which is compatible with S ;
- (ii) the ring $B_V(S)$ is noetherian;
- (iii) the conductor of $B_V(S)$ in $\mathbb{R}[V]$ is zero;
- (iv) S is Zariski dense in V at infinity.

We have (i) \Rightarrow (iii), (ii) \Rightarrow (iii) and (iii) \Leftrightarrow (iv).

Proof. (i) \Rightarrow (iii): Let $V \hookrightarrow X$ be a completion which is compatible with S , let Y be the union of the irreducible components Z of $X \setminus V$ with $Z(\mathbb{R}) = \emptyset$, and let $U = X \setminus Y$. By Thm. 3.8, the inclusion $V \hookrightarrow U$ induces an isomorphism $\mathcal{O}(U) \xrightarrow{\sim} B_V(S)$. Since S is unbounded we have $U \neq V$. By Lemma 5.6, $B_V(S)$ contains no non-zero ideal of $\mathbb{R}[V]$.

(ii) \Rightarrow (iii): We have $B \neq \mathbb{R}[V]$ since S is unbounded. Let J be an ideal of $\mathbb{R}[V]$ which is contained in $B_V(S)$. Since B is noetherian, Lemma 5.5 implies $J = (0)$.

The equivalence of (iii) and (iv) follows from Prop. 5.3. \square

Corollary 5.8. *Let V be a connected normal affine \mathbb{R} -variety, and let S be an unbounded semi-algebraic subset of $V(\mathbb{R})$. If S is not Zariski dense at infinity, then $B_V(S)$ is not noetherian.*

Proof. The vanishing ideal of the Zariski closure of S at infinity is a proper non-zero ideal of $\mathbb{R}[V]$ that is contained in $B_V(S)$. Hence the conductor of $B_V(S)$ in $\mathbb{R}[V]$ is non-zero, and so $B_V(S)$ is not noetherian by Prop. 5.7. \square

Remark 5.9. Condition (i) in Prop. 5.7 implies $B_V(S) \cong \mathcal{O}_U(U)$ for some \mathbb{R} -variety U (Thm. 3.8). In general, the ring $\mathcal{O}_U(U)$, and hence also $B_V(S)$, may fail to be noetherian, see 5.13 below. Therefore, (i) does not imply (ii) in Prop. 5.7.

The implication (ii) \Rightarrow (iv) in Prop. 5.7 can be strengthened as follows:

Proposition 5.10. *Let V be a normal affine \mathbb{R} -variety, and let S be a semi-algebraic subset of $V(\mathbb{R})$ for which the ring $B_V(S)$ is noetherian. Then for every semi-algebraic subset $T \subset S$ with $B_V(S) \neq B_V(T)$, the inequality*

$$\text{trdeg}_{\mathbb{R}} B_V(T) \leq \dim(S \setminus T) \tag{5.1}$$

holds.

Assume that S is unbounded and (for simplicity) closed, and that $B_V(S)$ is noetherian. Then conclusion (5.1) implies for any compact subset T of S that $S \setminus T$ is Zariski dense in $V(\mathbb{R})$. This means that S is Zariski dense at infinity, and shows that the implication (ii) \Rightarrow (iv) of 5.7 is contained in Prop. 5.10.

Proof. Let Z be the Zariski closure of $S \setminus T$ in V , and let $I = I_Z$ be the vanishing ideal of Z . Then $B(T)/I \cap B(T) \subset \mathbb{R}[V]/I$, and so

$$\text{trdeg}_{\mathbb{R}} B(T)/I \cap B(T) \leq \text{trdeg}_{\mathbb{R}} \mathbb{R}[V]/I = \dim Z = \dim(S \setminus T).$$

If (5.1) were false we would conclude $I \cap B(T) \neq (0)$. Now the ideal $I \cap B(T)$ of $B(T)$ is contained in $B(S)$. This contradicts Lemma 5.5 since $B(S)$ is integrally closed in $B(T)$ (even in $\mathbb{R}[V]$). \square

We now consider algebraic surfaces. Theorem 5.12 below is one of our main results. It is based on the results of Sections 3 and 4 and on the following theorem:

Theorem 5.11 (Zariski). *Let W be a normal quasi-projective surface over a field k of characteristic zero. Then the ring $\mathcal{O}_W(W)$ is finitely generated as a k -algebra.*

See Zariski [22], remarks on Theorem 1 at the end of the article. An alternative proof in modern language was given in [9], based on ideas of Nagata.

Theorem 5.12. *Let V be a connected normal affine surface over \mathbb{R} . For every semi-algebraic subset S of $V(\mathbb{R})$ which is regular at infinity, the \mathbb{R} -algebra $B_V(S)$ is finitely generated.*

Proof. By Theorem 4.5, V has a projective completion $V \hookrightarrow X$ which is compatible with S . If Y is the union of all irreducible components of $X \setminus V$ without a real point, then $\mathcal{O}_X(X \setminus Y) \xrightarrow{\sim} B_V(S)$ by Theorem 3.8. This implies the assertion, since $\mathcal{O}_X(X \setminus Y)$ is finitely generated by Zariski's theorem 5.11. \square

5.13. It is folklore knowledge that Zariski's theorem 5.11 ceases to be true in dimensions larger than two. However, it seems not so easy to localize a reference for this fact in the published literature.

A construction of a quasi-projective non-singular threefold whose ring of global sections is not finitely generated was given by Vakil [21]: Take an elliptic curve E over a field k , and let L, L' be invertible sheaves on E such that $\deg(L') > 0$, $\deg(L) = 0$ and L is not torsion in $\text{Pic}(E)$. Let W be the total space of the vector bundle $L \oplus L'$ on E . Then the ring $\mathcal{O}_W(W)$ is not noetherian. By a variation of this construction one can also obtain another example which is even quasi-affine (see [21]).

Using this construction for $k = \mathbb{R}$ in a case where the elliptic curve E is real, we can apply Theorem 4.11 and conclude:

Corollary 5.14. *There exists a non-singular affine \mathbb{R} -variety V of dimension three and a regular and basic closed semi-algebraic subset S of $V(\mathbb{R})$ for which the ring $B_V(S)$ is not noetherian.* \square

5.15. From results of Kuroda [10], it is possible to deduce the existence of a rational, normal, quasi-affine threefold whose ring of regular functions is not finitely generated. This was pointed out to us by Sebastian Krug, thereby answering a question raised in an earlier version of this paper.

5.16. By Zariski's theorem, an example as in 5.13 cannot exist in dimension two. However, there are examples of (non-normal) irreducible quasi-projective surfaces W for which $\mathcal{O}_W(W)$ is not finitely generated. One such construction of a quasi-affine example is due to Nagata ([13], [14], see also [9] for a detailed account).

The construction of two-dimensional varieties whose ring of global sections is not finitely generated becomes much easier when we allow the variety to be reducible. Here is an example (again see [9] for details): In affine 3-space with coordinates (x, y, z) let $U := V(xy)$ (the transversal union of two planes) and $L := V(y, z)$ (a line in one of the planes which intersects the other plane transversely). Then $W := U \setminus L$ is a reducible quasi-affine variety of dimension two for which the ring $\mathcal{O}_W(W)$ is not noetherian.

To interpret this example in terms of bounded functions, let

$$V = U \setminus (U \cap V(z)) = W \setminus V(x, z),$$

an open affine subvariety of W , and write $L' = V(x, z)$. Let S_0 be a regular, compact and basic closed semi-algebraic subset of $U(\mathbb{R})$ which is disjoint from L and compatible with L' . For example, $S_0 = \{(0, y, z) : (y - 2)^2 + z^2 \leq 1\}$ will do. Then $S = S_0 \cap V(\mathbb{R})$ is basic closed in $V(\mathbb{R})$, and $B_V(S) = \mathcal{O}_W(W)$ is not noetherian.

REFERENCES

- [1] E. Becker: The real holomorphy ring and sums of $2n$ -th powers. In: *Real Algebraic Geometry and Quadratic Forms* (Rennes, 1981), Lect. Notes Math. **959**, Springer, Berlin, 1982, pp. 139–181.

- [2] E. Becker, V. Powers: Sums of powers in rings and the real holomorphy ring. *J. reine angew. Math.* **480**, 71–103 (1996).
- [3] J. Bochnak, M. Coste, M.-F. Roy: *Real Algebraic Geometry*. *Erg. Math. Grenzgeb.* (3) **36**, Springer, Berlin, 1998.
- [4] S. D. Cutkosky: *Resolution of Singularities*. *Grad. Studies Math.* **63**, AMS, Providence, RI, 2004.
- [5] D. Eisenbud. *Commutative Algebra with a View Toward Algebraic Geometry*, *Grad. Texts Math.* **150**, Springer, New York, 1995.
- [6] R. Hartshorne: *Algebraic Geometry*. *Grad. Texts Math.* **52**, Springer, New York, NY, 1977.
- [7] S. Iitaka: *Algebraic Geometry*. *Grad. Texts Math.* **76**, Springer, New York, 1982.
- [8] J.-P. Jouanolou: *Théorèmes de Bertini et applications*. *Prog. Math.* **42**, Birkhäuser, Boston, MA, 1983.
- [9] S. Krug: Der Ring der globalen regulären Funktionen einer algebraischen Varietät. Diplomarbeit, Konstanz 2008.
- [10] S. Kuroda: A counterexample to the fourteenth problem of Hilbert in dimension three. *Michigan Math. J.* **53** 123–132 (2005).
- [11] M. Marshall: Approximating positive polynomials using sums of squares. *Canad. Math. Bull.* **46**, 400–418 (2003).
- [12] J.-P. Monnier: Anneaux d’holomorphic et Positivstellensatz archimédiens. *Manuscr. math.* **97**, 269–302 (1998).
- [13] M. Nagata: Addition and corrections to my paper “A treatise on the 14-th problem of Hilbert”. *Mem. Coll. Sci. Univ. Kyoto Ser. A Math.* **30**, 197–200 (1957).
- [14] M. Nagata: Lectures on the fourteenth problem of Hilbert. TIFR, Bombay, 1965.
- [15] D. Plaumann: Sums of squares on reducible real curves. *Math. Z.* (2010, to appear).
- [16] M. Roggero: Sui sistemi lineari e il gruppo delle classi di divisori di una varietà reale. *Ann. Mat. Pura Appl.* (4) **135**, 349–362 (1984).
- [17] C. Scheiderer: *Real and Étale Cohomology*. *Lect. Notes Math.* **1588**, Springer, Berlin, 1994.
- [18] C. Scheiderer: Sums of squares of regular functions on real algebraic varieties. *Trans. Am. Math. Soc.* **352**, 1039–1069 (1999).
- [19] C. Scheiderer: Sums of squares on real algebraic curves. *Math. Z.* **245**, 725–760 (2003).
- [20] M. Schweighofer: Iterated rings of bounded elements and generalizations of Schmüdgen’s Positivstellensatz. *J. reine angew. Math.* **554**, 19–45 (2003).
- [21] R. Vakil: An example of a nice variety whose ring of global sections is not finitely generated. Manuscript (2000), math.stanford.edu/~vakil/files/nonfg.pdf
- [22] O. Zariski: Interprétations algébrique-géométriques du quatorzième problème de Hilbert. *Bull. Sci. Math.* (2) **78**, 155–168 (1954).

UNIVERSITÄT KONSTANZ, FACHBEREICH MATHEMATIK UND STATISTIK, 78457 KONSTANZ, GERMANY

E-mail address: daniel.plaumann@uni-konstanz.de, claus.scheiderer@uni-konstanz.de